

CLAIMS

1. A method of synthesis of a chemical compound having the formula A-B-C
where the A is a chemiluminescent moiety,
5 B is an energy acceptor moiety, and
C is a biologically active moiety
comprising the steps of
forming a benzophenone,
forming a diaryl ethylene, and
10 performing at least one of
 - (a) attaching a precursor to generate a phthalhydrazide such as phthalimide, aminophthalic acid diester, aminophthalic acid dihydrazide, aminophthalic anhydride and phthalhydrazide protected by a hydrolyzable group to form the precursor-ethylene conjugate, and condensing two ethylene-precursor conjugates to form a precursor-pentadiene conjugate, and
 - 15 (b) condensing two diaryl ethylene to form a pentadiene, and attaching a precursor to generate a phthalhydrazide such as phthalimide, aminophthalic acid diester, aminophthalic acid dihydrazide, aminophthalic anhydride and phthalhydrazide protected by a hydrolyzable group, to form the precursor-pentadiene conjugate, and
converting the precursor to the phthalhydrazide by at least one of the corresponding
20 reactions
 - phthalimide with hydrazine,
 - aminophthalic acid diester with hydrazine,
 - aminophthalic anhydride with hydrazine, and
 - hydrolysis of phthalhydrazide protected by a hydrolyzable group to form a carrier
25 compound, and
reacting the carrier compound with the biologically active moiety to form a corresponding conjugate.
2. The method of synthesis of the compound of claim 1 wherein the compound serves to
30 delivery the C moiety to a desired biological compartment.
3. The method of synthesis of the compound of claim 1 wherein the compound is a prodrug.
4. The method of synthesis of the compound of claim 3 wherein the compound serves as a
35 prodrug for at least one of antiviral agents for the treatment of viral infections and anticancer agents for the treatment of cancers.
5. The method of synthesis of the compound of claim 4 wherein the compound serves as a prodrug for the treatment of at least one of the group of viruses comprising Human

Immunodeficiency Virus (HIV), herpes viruses such as Herpes Simplex Virus, (HSV), Epstein-Barr Virus (EBV), Varicella Zoster (VZV), Cytomegalovirus (CMV), HSV-6, and HSV-8 (Kaposi's sarcoma), Human Papilloma Virus (HPV), rhinoviruses, and hepatitis-linked viruses.

- 5 6. The method of synthesis of the compound of claim 4 wherein the compound serves as a prodrug for the treatment of at least one of the group of cancers comprising colon, breast, lung, renal, retinal, and skin.
7. The method of synthesis of the compound of claim 3 wherein the prodrugs have increased
10 bioavailability.
8. The method of synthesis of the compound of claim 2 wherein the compound is a cellular permeant prodrug.
- 15 9. The method of synthesis of the compound of claim 8 wherein intracellular drug release occurs when the prodrug reacts with cellular free radicals via a mechanism involving chemiluminescence, photochromism, and intramolecular energy transfer.
10. The method of synthesis of the compound of claim 1 wherein the C moiety is a
20 pharmaceutical agent or drug.
11. The method of synthesis of the compound of claim 10 wherein the pharmaceutical agent is at least one of the group of antilipidemic drugs, anticholesterol drugs, contraceptive agents, anticoagulants, anti-inflammatory agents, immuno-suppressive drugs, antiarrhythmic agents,
25 antineoplastic drugs, antihypertensive drugs, epinephrine blocking agents, cardiac inotropic drugs, antidepressant drugs, diuretics, antifungal agents, antibacterial drugs, anxiolytic agents, sedatives, muscle relaxants, anticonvulsants, agents for the treatment of ulcer disease, agents for the treatment of asthma and hypersensitivity reactions, antithromboembolic agents, agents for the treatment of muscular dystrophy, agents to effect a therapeutic abortion, agents for the treatment
30 of anemia, agents to improve allograft survival, agents for the treatment of disorders of purine metabolism, agents for the treatment of ischemic heart disease, agents for the treatment of opiate withdrawal, agents which activate the effects of secondary messenger inositol triphosphate, agents to block spinal reflexes, and antiviral agents including a drug for the treatment of AIDS.
- 35 12. The method of synthesis of the compound of claim 1 wherein the C moiety is released by an oxidation reduction reaction with the target cell's electron carriers or by reaction with free radicals produced as a consequence of electron transport.
13. The method of synthesis of the compound of claim 12 wherein the C moiety is released

into a desired compartment in active form.

14. The method of synthesis of the compound of claim 13 wherein the released C moiety has a greater therapeutic effect or therapeutic ratio relative to the free C agent alone.

5

15. The method of synthesis of the compound of claim 14 wherein the released C moiety has a greater therapeutic effect or therapeutic ratio relative to the free C agent alone as a consequence of at least one of altered pharmacokinetics or pharmacodynamics such as a desirable kinetics of release, a resistance to inactivation or excretion, greater solubility, enhanced absorption, a
10 diminished toxicity, or greater access to the cellular or biological compartment which is the site of action of C.

16. The method of synthesis of the compound of claim 1 wherein A represents a functionality which undergoes at least one of

15 an oxidation reduction reaction where electrons are transferred directly between A and the target cell's electron carriers, and

a reaction with free radicals of oxygen which are produced as a consequence of electron transport

20 such that an excited state is produced in A as a consequence of its participation in one of these reactions.

17. The method of synthesis of the compound of claim 16 wherein A undergoes intramolecular energy transfer from its own excited state to the B functionality which is an energy acceptor.

25

18. The method of synthesis of the compound of claim 17 wherein upon receiving energy from A, B achieves an excited state which relaxes through heterolytic cleavage of the covalent bond of B with C where C is a drug moiety which is released into the environment.

30 19. The method of synthesis of the compound of claim 18 wherein the released drug molecule effects a therapeutic functional change by a mechanism which comprises receptor mediated mechanisms including reversible and irreversible competitive agonism or antagonism including a molecule known as a suicide substrate or a transition state analogue mechanism or a noncompetitive or uncompetitive agonism or antagonism or the action is by a nonreceptor
35 mediated mechanism including a "counterfeit incorporation-mechanism".

20. The method of synthesis of the compound of claim 1 wherein the chemiluminescent molecule comprises at least one of the group of

molecules undergoing reaction involving peroxides and oxygen free radicals,

molecules undergoing reaction involving oxidation or reduction, and
molecules undergoing both reaction with peroxides and oxygen free radicals followed by
an oxidation or reduction reaction.

5 21. The method of synthesis of the compound of claim 20 wherein the chemiluminescent
molecule comprises at least one of the group of luminol and its derivatives, lucigenin and its
derivatives, Lophine and its derivatives, acridinium esters and acridans, tetraphenylpyrrole,
phthalhydrazides, acyloins, biacridinium salts, vinylcarbonyls, vinylnitriles, tetrakis
(dimethylamino) ethylene, acylperoxides, indoles, tetracarbazoles and active oxalates.

10

22. The method of synthesis of the compound of claim 20 wherein the chemiluminescent
molecule comprises at least one of the group of ruthenium chelates 2, 6-diaminopyrene, or cation
radicals and molecules which follow a Chemically Initiated Electron Exchange Luminescence
mechanism such as certain dioxetans and dioxetanones.

15


23. The method of synthesis of the compound of claim 20 wherein the chemiluminescent
molecule comprises at least one of the group of dioxene derivatives and other compounds that
form a dioxetan by reaction with superoxide and then produce efficient chemiluminescence by a
CIEEL mechanism.

20


24. The method of synthesis of the compound of claim 20 wherein the chemiluminescent
molecule comprises at least one of the group of

25


2,6-diaminopyrene




Aminophthalhydrazide




Dioxene




Imidazole derivatives




Sulfonyloxamides




Indole derivatives




Tetrakis(dialkylamino)-ethylene




2,5,7,8-tetraoxabicyclo-[4.2.0.]octane




Dioxetan




Lucigenin




Lophine




Acridinium esters




Active oxalate




Tris-2,2'-bipyridinedi-chlororuthenium (II)



Dioxetanone



Dipheyl peroxide



5

25. The method of synthesis of the compound of claim 1 wherein the B moiety is a photochromic compound.

10

26. The method of synthesis of the compound of claim 25 wherein the photochromic compound comprises one which demonstrate photochromic behavior with electromagnetic radiation and bleaching agents.

27. The method of synthesis of the compound of claim 26 wherein the A functionality is chemiluminescent, and the B functionality is such that the photodissociative drug release spectrum of B overlaps the chemiluminescence spectrum of A.

5 28. The method of synthesis of the compound of claim 25 wherein the photochromic compound comprises a cationic dye.

29. The method of synthesis of the compound of claim 28 wherein the cationic dye comprises at least one of a di and triarylmethane dyes, triarylmethane lactones and cyclic ether dyes,
10 cationic indoles, pyronines, phthaleins, oxazines, thiazines, acridines, phenazines, and anthocyanidins, and cationic polymethine dyes and azo and diazopolymethines, styryls, cyanines, hemicyanines, dialkylaminopolyenes, and other related dyes.


30. The method of synthesis of the compound of claim 28 wherein the cationic dye comprises
15 at least one of

	Malachite Green	42000
	Helvetia Green	42020
20	Basic Blue 1	42025
	Brilliant Blue	
	Setoglaurine	
	Basic Green 1	42040
	Brilliant Green	
25	Acid Blue 1	42045
	Xylene Blue VS	
	Patent Blue V	
	Alphazurine 2G	
	Acid Blue 3	42051
30	Brilliant Blue V	
	Patent Blue V	
	Food Green 3	42053
	FDC Green 3	
	Acid Green 6	42075
35	Light Green SF Bluish	
	Acid Blue 7	42080
	Xylene Blue AS	
	Patent Blue A	
	Acid Green 3	42085


	Acid Blue 9	42090
	Erioglaucine	
	Acid Green 5	42095
	Light Green SF Yellowish	
5	Acid Green 9	42100
	Erioviridene B	
	Acid Blue 147	42135
	Xylene Cyanol FF	
	Basic Red 9	42500
10	Pararosaniline	
	Basic Violet 14	42510
	Fuchsin	
	Magenta	
	Basic Fuchsin	42510B
15	Basic Violet 2	42520
	New Magenta	
	Hoffman Violet	42530
	Iodine Violet	
	Basic Violet 1	42535
20	Methyl Violet	
	Basic Violet 13	42536
	Methyl Violet 6B	
	Basic Violet 3	42555
	Crystal Violet	
25	Gentian Violet	
	Iodine Green	42556
	Basic Blue 8	42563
	Victoria Blue 4R	
	Acid Blue 13	42571
30	Fast Acid Violet 10B	
	Acid Blue 75	42576
	Eriocyanine A	
	Methyl Green	42585
	Ethyl Green	42590
35	Basic Violet 4	42600
	Ethyl Violet	
	Acid Violet 49	42640
	Wool Violet 5BN	
	Acid Blue 15	42645

	Brilliant Milling Blue B	
	Acid Violet 17	42650
	Acid Violet 6B	
	Wood Violet 4BN	
5	Formyl Violet	
	Acid Violet 5BS Conc.	
	Acid Violet 19	42685
	Acid Fuchsin	
	Red Violet 5R	42690
10	Acid Blue 22	42755
	Aniline Blue	
	Soluble Blue	
	Solvent Blue 3	42775
	Solvent Blue 3	42780
15	Methyl Blue	
	Aurin	43800
	Mordant Blue 3	43820
	Eriochrome Cyanine R	
	Acid Green 16	44025
20	Naphthalene Green V	
	Pontacyl Green NV Extra	
	Basic Blue 11	44040
	Victoria Blue R	
	Basic Blue 15	44085
25	Night Blue	
	Acid Green 50	44090
	Wool Green S	
	Kiton Green S. Conc.	
	Basic Green 3	
30	Sevron Green B	
	Brilliant Blue F & R Extra	
	Brilliant Green Sulfonate	
	Hexakis (hydroxyethyl)	


Pararosaniline




New Green




Phenolphthalein




Malachite Green Ethiodide




Hydroxyalkylated Pararosanilines




Hydroxyalkylated New Fuchsins




New Yellow




Doebner's Violet




New Red




Bis(hydroxyethyl) Doebner's Violet




"New Magenta"










Tetrakis(hydroxyethyl) Doebner's Violet




Trichloro Crystal Violet



Slow Red





5 ^a Only the cyanide, bisulfite, and hydroxide ions are considered, regardless of the other anions present in the solution.

^b More detailed descriptions of the compositions of photochromic materials tested are given in Macnair's review [255; tables 1A-4].

^c Ethanol.

10 ^d Diethyl ether.

^e 1,2-Dichloroethane.

^f 1,1-Dichloroethane, cyclohexane-1,1-dichloroethane, or cyclohexane-1,2-dichloroethane mixtures.

^g Benzene.

^h Dimethylsulfoxide, neat and aqueous.

15 ⁱ Acetone.

^j ACETIC ACID.

^k Ethyl acetate.

- ^l Ethyl bromide.
- ^m 2-Methoxyethanol.
- ⁿ Chloroform.
- ^o Ethanol with KCN.
- 5 ^p Ethanol with KOH.
- ^q Carboxylic acids-acetic to stearic; hydrocinnamic acid; ethyl and butyl acid phthalates.
- ^r Octadecylnitrile, tributyl phosphate, aniline, 2-(p-tert-butylphenoxy)ethanol, tetraethyleneglycol dimethyl ether, or poly(ethylene glycols).
- ^s Amides-formamide to stearamide; methylformamide or methylacetamide; dimethyl- or diethyl-
- 10 formamide or acetamide.
- ^t Three-to-one solutions of cellulose acetate with any of the following five-to-one plasticizer mixtures: butyl stearate, Polyethylene Glycol 600-butyl acetoxysearate, butyl stearate, or Dowanol EP-butyl acetoxysearate.
- ^u Water containing SO₂.
- 15 ^v Water containing bisulfite and papain.
- ^w Poly(vinyl alcohol) with dimethylsulfoxide (5:1).
- ^x Films, containing residual solvent, cast from the following solutions: ethanol-acetone solutions of vinyl acetate-vinyl alcohol copolymer; aqueous poly(vinyl alcohol); aqueous poly(vinyl pyrrolidone); or aqueous methyl vinylether-maleic acid copolymer.
- 20 ^y Methanol-dioxane with aqueous NH₄ HSO₃.
- ^z Paper impregnated with a toluene solution of poly(methyl methacrylate), stearic acid, and 2-(p-tert-butylphenoxy)ethanol, then dried.
- ^{aa} Intramicellar impregnation of cellulose with the following swelling agents: n-propylamine, n-butylamine, n-hexylamine, 2-aminoethanol, dimethylformamide, acetic acid, dimethylsulfoxide,
- 25 methylacetamide, dimethylacetamide, or formamide.
- ^{bb} Films cast from an approximately 4:3 mixture of a 20% solution and cellulose acetate butyrate in toluene-ethyl acetate (1:1) and triallycyanurate in dioxane.
- cc FILMS CAST FROM A 2:1 MIXTURE OF A 25% SOLUTION OF CELLULOSE ACETATE BUTYRATE IN TOLUENE-ETHYL ACETATE (1:1) AND THE TITANIUM ESTERS OF N,N,N', N'-TETRAKIS(2-HYDROXYPROPYL) ETHYLENEDIAMINE.**
- 30 ^{dd} Pure water.
- ^{ee} Films cast from aqueous gelatin or other hydrocolloids.
- ^{ff} Dimethylsulfoxide with methanolic KCN.
- ^{gg} 2-Methoxyethanol with methanolic KCN.
- 35 ^{hh} Water or aqueous methanol containing bisulfite.
- ⁱⁱ Paper impregnated with m-dimethoxybenzene, acetonitrile, acetic acid, or phenyl methyl carbinol.
- ^{jj} Ethanol-benzene.
- ^{kk} Aqueous ethanol, methanol, aqueous methanol, aqueous acetone, benzene-methanol, carbon tetrachloride-methanol, cyclohexane-methanol, or chloroform-methanol.

^{ll} Films cast from 3:1 solutions of cellulose acetate and either Polyethylene Glycol 600 .RTM. or ethylene glycol phenyl ether as plasticizer.

^{mmm} Films, containing residual solvent, cast from solutions of either cellulose acetate in 2-methoxyethanol or poly(vinyl alcohol) in aqueous ethanol.

5 ⁿⁿ Films, containing residual solvent, cast from solutions of either cellulose acetate butyrate in 2-methoxyethanol or poly(vinyl acetate) in methanol.

^{oo} Ethanol containing ammonia.

^{pp} Aqueous methanol containing NH_4HSO_3 and urease.

^{qq} Aqueous methanol containing NH_4HSO_3 , with or without sodium dithionite.

10 ^{rr} Aqueous acid at pH 1.

^{ss} Aqueous ammonia containing KCN.


^{tt} Paper impregnated with aqueous solutions with or without hydrocolloids.





^{uu} 2-Methoxyethanol containing HCl.

^{vv} Aqueous methanol containing NH_4HSO_3 , and glucose oxidase.

15 ^{ww} 9:1 Methanol-water.


^{xx} Aqueous NaOH.





Photochromic Polymethine Dyes

5




Ar

n

10	C ₆ H ₅	0, 1, 2
	4-(CH ₃) ₂ NC ₆ H ₄	0, 1, 2
	4-(CH ₃) ₂ CHC ₆ H ₄	0, 1, 2, 3, 4
	4-CH ₃ OC ₆ H ₄	0, 1, 2
	4-C ₄ H ₉ OC ₆ H ₄	0, 1, 2
15	3-CH ₃ C ₆ H ₄	1, 2
	4-t-C ₄ H ₉ C ₆ H ₄	1, 2



	4-C ₂ H ₅ OC ₆ H ₄	1, 2
	4-C ₅ H ₁₁ C ₆ H ₄	1, 2
	4-FC ₆ H ₄	1
	4-Fsub ₃ CC ₆ H ₄	1
5	2-(C ₆ H ₅) ₂ NC ₆ H ₄	1
	3,4-H ₂ N(OCH ₃)C ₆ H ₃	1
	2-Naphthyl	1, 2
	4-ClC ₆ H ₄	2
	2,4-Cl ₂ C ₆ H ₃	2
10	1-Naphthyl	2

















R

R

15








20












Miscellaneous polyenes





5





























Salt-isomerism type phototropic dyes

5


Night Blue




Victoria Blue R




Brilliant Milling Blue B
Brilliant Blue F & R Ex.
Eriocyanine A




Methyl Blue




Aniline Blue




Eriochrome Cyanine R




Methyl Violet 6B




Iodine Green




Aniline Blue




Wool Violet 5 BN




Wool Violet 4 EM




Light Green SF
Yellowish




Iodine Violet




Methyl Violet




Crystal Violet




Red-Violet 5R




Brilliant Green "B"




Di-[4(N,N-diethylamine)phenyl]-[4(N,N-diethylamine-2-methyl) phenyl] methyl carbonium




Tri-[4(N,N-dipropylamino)phenyl] methyl carbonium




Di-[4(N,N-diethylamino)phenyl]-[4(ethylamino)-phenyl] methyl carbonium




Di-[4(N,N-diethylamino)phenyl]-[4(N,N-diethylamino)naphthyl] methyl carbonium



Di-[4(N,N-dimethylamino)phenyl]-[4(hydroxy)phenyl]
methyl carbonium



Tri-[4(N-propylamino)phenyl] methyl carbonium




Hectolene Blue DS-1398


Hectolene Blue DS-1823

Sevron Brilliant Red 4G

Di-[4(N,N-dimethylamino)phenyl]-[4(hydroxy)phenyl]
methyl carbonium



Tri-[4(N-propylamino)phenyl] methyl carbonium



Hectolene Blue DS-1398

Hectolene Blue DS-1823

Sevron Brilliant Red 4G


Genacryl Red 6B

Genacryl Pink G


Sevron Brilliant - Red B

Sevron Brilliant - Red 3B


1,5-bis-[4(N,N-dimethylamino)phenyl]-1,5-bis-
(phenyl)divinyl carbonium trifluoroacetate




1,1,3,3-tetrakis[4(N,N-dimethylamino)phenyl]
vinyl carbonium perchlorate




1,5-bis-[4(N,N-dimethylamino)phenyl]-1,5-bis-
(phenyl) divinyl carbonium p-toluenesulfonate




1,7-bis[4(N,N-dimethylamino)phenyl]-1,7-bis-
(2,4-dichlorophenyl) trivinyl carbonium perchlorate




Di-[4(N,N-dimethylamino)phenyl vinyl]-[2,4-di-
phenyl-6-methane thiopyran] methyl carbonium perchlorate




1,7-bis-[4(N,N-dimethylamino)phenyl]-1,7-bis-
(4-chlorophenyl) trivinyl carbonium trifluoroacetate




1,1,3-tris-[4-(*N,N*-dimethylamino)phenyl] divinyl
carbonium perchlorate




1,1,7,7-tetrakis-[4-(*N,N*-dimethylamino)phenyl]
trivinyl carbonium perchlorate




1,3-bis-[4-(*N,N*-dimethylamino)phenyl]-1,3-bis-
(phenyl) vinyl carbonium perchlorate




1,1,5,5-tetrakis-[4-(*N,N*-dimethylamino)phenyl]
divinyl carbonium perchlorate



1,5-bis-[4-(*N,N*-dimethylamino)phenyl]-1,5-bis-
(phenyl) divinyl carbonium perchlorate




1,7-bis-[4-(N,N-dimethylamino)phenyl]-1,7-bis-(phenyl) trivinyl carbonium trifluoroacetate




CF₃COO—

1(1,3,3-trimethyl indoline)-2-[4-(N,N-dimethylamino)phenyl] ethylene carbonium perchlorate




ClO₄—

1(1,3,3-trimethyl indoline)-4-[4-(N,N-dimethylamino)phenyl] butylene carbonium perchlorate




ClO₄—

1,1,3,3-tetrakis-[4-(N,N-diethylamino)phenyl] vinyl carbonium perchlorate




ClO₄—

1,1-bis-[4-(N,N-diethylamino)phenyl]-3,3-bis-[4-(N,N-dimethylamino)phenyl] vinyl carbonium perchlorate




ClO₄—


1,1,5,5-tetrakis-[4-(N,N-diethylamino)phenyl]
divinyl carbonium perchlorate




1,1-bis-[4-(N,N-dimethylamino)phenyl]-3-[4-(amino)
phenyl]-3-methylvinyl carbonium perchlorate




Tris-[1,1-bis-[4-(N,N-dimethylamino)phenyl]
ethylene] methyl carbonium perchlorate




Tris-[1,1-bis-[4-(N,N-diethylamino)phenyl]
ethylene] methyl carbonium perchlorate




1,1,5-tris-[4-(N,N-dimethylamino)phenyl] divinyl
carbonium perchlorate




N[4-(N,N-dimethylamino) cinnamylidene] auramine




1,1-bis-[4-(N,N-dimethylamino)phenyl]-3,4-bis-
(phenyl)]-3,4-diazo butene carbonium



1,1,5,5-tetrakis-[4-(N,N-dimethylamino)phenyl]-
2,3-diazo pentene carbonium





N-(N',N'-dimethylamino cinnamylidene)-N,N-diphenyl
ammonium



Azo Polymethines

Dyes of the general structural type





5

31. The method of synthesis of the compound of claim 10 wherein the C moiety is any molecule which exhibits bleaching behavior with the B moiety and has an increased therapeutic effect or therapeutic ratio as a consequence of its delivery as part of a prodrug.

10 32. The method of synthesis of the compound of claim 29 wherein the C moiety has a nucleophilic group that bonds to the B moiety.


33. The method of synthesis of the compound of claim 32 wherein the C moiety is derivatized to have a nucleophilic group that bonds to the B moiety.

15


34. The method of synthesis of the compound of claim 33 wherein the C moiety is derivatized by at least one of the nucleophilic groups comprising cinnamate, sulfite, phosphate, carboxylate, thiol, amide, alkoxide, or amine.

35. The method of synthesis of the compound of claim 10 wherein the C moiety is at least one of the group of


Captopril




Prostaglandin E₂




2,3-dichloro- α -methylbenzylamine




3'-deoxy-S-adenosyl-L-homocysteine




Sinefungin




3,5-diiodo-4-hydroxybenzoic acid




6,6'-dithiobis (9-B-D-ribofuranosylpurine)




γ -aminobutyric acid




Gabaculine




N-(5'-phosphopyridoxy)-4-aminobutyric acid




4-amino-hex-5-enoic acid




Baclofen




Adenosine




3-hydroxy-3-methylglutarate




Campactin





But-3-ynoyl-CoA




Suramin




L-3-iodotyrosine

L-3-iodo- α -methyltyrosine


Disodium cromoglycate




Adenosine
3',5'-cyclic monophosphate




D,L-B-(5-hydroxy-3-indolyl)- α -hydrazino- α -propionic acid




D,L- α -hydrazino- α -methyl-dopa




α -methyl-dopa




5-(3,4-dihydroxycinnamoyl)salicylic acid




N-(phosphonacetyl)-L-aspartate




P-glycolohydroxamate




5-(p-sulfamylphenyl)-azosalicylic acid




Coformycin



Formycin B




Thioinosinate




Phosphonoformate




Phosphonoacetate




Ridavirin




Sotalol




Cimetidine




Fuscaric acid




2-mercaptoethylamine




Mimosine




U-7130




Iproniazid




Trans-4-aminocrotonic acid




NSD 1055



Nicotinic acid



Kynaurenic acid



NC1=NC=NC2=C1N=CN2C(CO)COC=C(C(=O)O)C=CC(=O)NNC(=O)OCC(O)C(=O)C(=O)N[C@@H]1[C@H](O)[C@@H](O)[C@H](O)[C@@H]1N2C(=O)N=C(C(=O)O)C(=O)N2R1NC(=O)C1CN2C(=C(C(=O)O)C(S)C2)R2CC(C)S[C@H]1C(=O)NC(=O)C1

36. The method of synthesis of the compound of claim 10 wherein the C moiety is at least one or a derivative or analog of one of the group of

10 prostaglandin A.sub.1 A.sub.2 B.sub.1 E.sub.1 , E.sub.2 or an analog which possesses a
vasodilatory effect on coronary arteries and other human vascular beds .

prostaglandin A, E, or an analogue prostaglandin which possesses natriuretic and diuretic

prostaglandin E.sub.1, E.sub.2 or an analogue which causes bronchial dilatation

prostaglandin F2 or an analogue which causes abortion by luteolysis
prostaglandin A.sub.2, E.sub.1, E.sub.2, or an analogue which induces erythropoiesis
prostaglandin E or an analogue which modulates T lymphocytes to decrease their ability
to reject an allogenic graft

5 2'-isopropyl-4'-(trimethylammonium chloride)-5'-methylphenyl piperidine -1-carboxylate
(Amo 1618) or an analog which inhibits the cyclization of trans-geranyl-geranyl-PP to copalyl-
PP during Kaurene synthesis

adenosine cyclic 3', 5'-monophosphate or an analogue which inhibits the release and
formation of phlogistic mediators such as histamine and kinins

10 4'-sulfamylphenyl

2-azo -7-acetamid-1-hydroxynaphthalene-3,6-disulfonate (Neoprontosil), 4'-sulfamyl-2,
4-diaminoazobenzene (Prontosil), or 5-(p-sulfamylphenylazo) salicylic acid (Lutazol) or analog
which possess potent carbonic acid anhydrase inhibition

analogue of S-adenosyl homocysteine or sinefungin

15 phosphoglycolohydroxamate which inhibits Class II aldolases present in bacterial and
fungi and is noninhibitory of Class I aldolases present in animals,

inosine analogue such as formycin B which inhibits nucleotide phosphorylase during
nucleotide metabolism

phosphonoformate (Foscarnet) or an analog which inhibits the HIV reverse transcriptase
20 enzyme

gamma.-amino-butyric acid (GABA) or an analog which is the major inhibitory
neurotransmitter in the mammalian central nervous system

gabaculine, N-(5'-phosphopyridoxyl)-4-aminobutyric acid, ethanolamine -o-sulfate,
.gamma.-vinyl GABA, or .gamma.-acetylenic GABA or an analog that is an inhibitor of the

25 GABA-degrading enzyme, GABA: 2-oxoglutarate aminotransferase

Baclofen or a compound that inhibits GABA release

an oligonucleotide which binds to RNA or DNA and blocks transcription or translation of
HIV or P-glycoprotein gene products adenosine which binds to brain purinergic receptors to
suppress opiate withdrawal

30 adenosine which causes coronary vasodilatation

3-hydroxy-3-methylglutarate, 3-hydroxybutyrate, 3-hydroxy-3-methylpentanoate, 4-
bromocrotonyl-CoA, but-3-ynoyl-CoA, pent-3-ynoyl-CoA, dec-3-ynoyl-CoA, ML-236A, ML-
236B (compactin), ML-236C, mevinolin, mevinolinic acid, or a mevalonic acid analogue which
is an inhibitor of 3-hydroxy-3-methylglutaryl-CoA reductase which catalyzes the rate-limiting
35 and irreversible step of cholesterol synthesis where inhibition at this step does not lead to the
accumulation of nonmetabolizable precursors

thioinosinate which suppresses T lymphocytes

Suramin, which is a powerful inhibitor of energy driven calcium uptake by the
sarcoplasmic reticulum and is an intracellular inhibitor of Na.sup.+ -K.sup.+ ATPase where both

activities increase intracellular calcium concentrations with a concomitant inotropic effect

norepinephrine N-methyltransferase inhibitor such as 2,3-dichloro- α -methylbenzylamine, 2,3-dichlorobenzylamine, 2,3-dichlorobenzamidine, or 3,4-dichlorophenylacetamidine

5 adenosine cyclic 3', 5'-monophosphate or a cAMP analogue which blocks the synthesis of fatty acids and cholesterol in the liver is an antilipidemic agent,

an inhibitor of dihydroxyphenylalanine decarboxylase during the synthesis of epinephrine and norepinephrine such as psitectorigenin, genistein, 3', 4',5,7-tetrahydroxy-8-methylisoflavone, orbol, 8-hydroxygenistein, 3',5,7-trihydroxy-4',6-dimethylisoflavone, 3',5,7-trihydroxy-4',8-
10 dimethoxyisoflavone, D,L-B-(5-hydroxy-3-indolyl)- α -hydrazinopropionic acid, D,L- α -hydrazino- α -methyl-dopa, D,L-B-(3-indolyl)- α -hydrazinopropionic acid, a derivative of phenylalanine such as N-methyl-3,4-dopa, α -acetamido-3,4-dimethoxycinnamic acid, DL- α -methyl-3,4-dopa, α -methyl-B-(3-hydroxy-4-methoxyphenyl)alanine, α -methyl-3,4-dimethoxyphenylalanine, or d-catechin; D,L-B-(3-indolyl)- α -methyl- α -
15 hydrazinopropionic acid (R)-3 β ,4-dihydroxyphenyl-1-fluoropropylamine, (S)- α -fluoromethyl-dopa, (S)- α -fluoromethyltyrosine, 5-(3,4-dihydroxycinnamoyl) salicylic acid, 3-hydroxycinnamic acid, caffeic acid, 3-mercaptopropionic acid, α -methyl-3-hydroxycinnamic acid, α -ethyl-3-hydroxycinnamic acid, 3-hydroxy-*w*-nitrostyrene, 3,4-dihydroxyhydrocinnamic acid, 3-hydroxybenzalacetone, 3-hydroxychalcone, 3-hydroxybenzal
20 furanyl ketone, 3-hydroxybenzal thiophenyl ketone, 3',4'-dihydroxyflavone, 8-O-glucoseflavone, flavone, 3-hydroxyphenyl pyruvic acid, 3,4-dihydroxyphenylpyruvic acid phenylthiopyruvic acid, 4-hydroxyphenylpyruvic acid, dithiosalicylic acid, 1-hydroxy-2-naphthoic acid, 3-hydroxy-7-sulfo-2-naphthoic acid, 3,5-dihydroxy-2-naphthoic acid, 4-chlorocinnamic acid, 2-chlorocinnamic acid, 2,4-dichlorocinnamic acid, 3-nitrocinnamic acid, 3,5-dibromo-2-
25 hydroxycinnamic acid, 2,4,6-triiodo-3-hydroxycinnamic acid, 2-hydroxy-4'-cyanochalcone, 4-(4-hydroxycinnamoyl) benzylnitrile, 2-(4-hydroxycinnamoyl)-1,4-dihydroxybenzene, quercetin-6'-sulfonic acid, 5-(2-hydroxy-3,5-dibromocinnamoyl) salicylic acid or 5-(3-hydroxycinnamoyl) salicylic acid

an inhibitor of acrosin, a proteolytic enzyme located in the acrosome of sperm, such as
30 tosyl lysine chloromethyl ketone, N- α -tosyl-L-arginine chloromethyl ketone, or ethyl p-guanidinobenzoate,

adenosine cyclic 3',5'-monophosphate (cAMP), N^{sup.6}, O^{sup.2}-dibutyryl-adenosine cyclic 3',5'-monophosphate or an analogue which produces an inotropic response,

adenosine kinase enzyme inhibitor such as 6,6'-dithiobis (9-B-D-ribofuranosyl)purine),

35 inhibitor of monoamine oxidase such as phenylhydrazine, phenylethylidenehydrazine, isopropylhydrazine, or iproniazid,

an inhibitor of catechol-O-methyltransferase such as 3,5-diiodo-4-hydroxybenzoic acid, S-3'-deoxyadenosyl-L-homocysteine, pyrogallol, R04-4602, gallic acid, 3,5-dihydroxy-4-methylbenzoic acid, 1,3-dihydroxy-2-methoxybenzene, 1-hydroxy-2,3-dimethoxybenzene, 2-

hydroxy-1,3-dimethoxybenzene, 1,3-dihydroxy-4-methoxybenzene, catechol, 3,4-dihydroxybenzoic acid, caffeic acid, 5,6-dihydroxyindole, noradrenaline, dopacetamide, H 22/54, quercetin, nordihydroguaiaretic acid, U-0521, arterenone, methylspinazarin, MK 486, dopa, papaveroline, isoprenaline, 7,8-dihydroxy-chlorpromazine, 3-hydroxy-4-pyridone, tetrahydroisoquinoline pyridoxal 5'-phosphate, iodoacetic acid, 3-mercaptoptyramine, dehydrodicaffeic acid dilactone, methylspinazarin, 3',5,7-trihydroxy-4',6-dimethoxyisoflavone, 3',5,7-trihydroxy-4',8-dimethoxyisoflavone, 6,7-dihydromethylspinazarin, S-adenosylhomocysteine, S-tubercidinylhomocysteine, 3',8-dihydroxy-4',6,7-trimethoxyisoflavone, 7-O-methylspinachrome B, 6-(3-hydroxybutyl)-7-O-methylspinachrome B, 3,5-diiodosalicylic acid, or pyridoxal-5'-phosphate,

an inhibitor of adenosine deaminase which blocks the metabolism of adenosine such as coformycin, arabinosyl-6-thiopurine, 6-methylthioinosine, 6-thioinosine, 6-thioguanosine, N^{sup.1}-methyladenosine, N^{sup.6}-methyladenosine, 2-fluorodeoxyadenosine, 2-fluoroadenosine, inosine, 2'-deoxyinosine, deoxycostomycin, 1,6-dihydro-6-hydroxymethyl purine ribonucleoside, erythro-9-(2-hydroxy-3-nonyl)adenine, or 9-B-D-arabinofuranosyl-6-hydroxylaminopurine,

an inhibitor of adenylate kinase, 5'-nucleotidase, and adenosine translocase such as p^{sup.1} p^{sup.5}-diadenosine pentaphosphate, α -, β -methylene adenosine diphosphate, and nitrobenzyl-6-thioinosine, respectively,

an inhibitor of γ -aminobutyric acid uptake such as D,L-2,4-diaminobutyric acid, D,L-B-hydroxy GABA, (-)-nipecotic acid, trans-4-aminocrotonic acid, cis-3-aminocyclopentane-1-carboxylic acid, trans-3-aminocyclopentane-1-carboxylic acid, β -guanidinopropionic acid, homohypotaurine, 4-aminopentanoic acid, homotaurine, β -alanine, imidazoleacetic acid, 6-aminohexanoic acid, D,L-carnitine, D,L-2,6-diaminopimelic acid, D,L-2-fluoro GABA, guanidino acetic acid, 2-hydrazinopropionic acid, taurine, D,L-ornithine, or sulphanilamine which potentiates the inhibitory action of GABA,

inositol 1,4,5-triphosphate,

guanosine 5' cyclic monophosphate or 8-bromo guanosine 5' cyclic monophosphate which relaxes smooth muscle,

an inhibitor of the uptake system for glycine, the inhibitory synaptic transmitter of the spinal cord, such as hydrazinoacetic acid,

isoquinoline-sulfonamide inhibitor of protein kinase C, cAMP-dependant protein kinase, or cGMP-dependent protein kinase such as N-(2-aminoethyl)-5-isoquinolinesulfonamide,

Ribavirin which is active against HSV-1 and 2, hepatitis, and influenza viruses, or phosphonoacetic acid which is a highly specific inhibitor of Herpes Simplex virus induced polymerase and is active against HSV-1 and HSV-2, or adenine arabinoside (ara-A), cytosine arabinoside (Ara-C), ara-A 5'-monophosphate (ara-AMP), or hypoxanthine arabinoside (ara-Hx) which is active against HSV or phagocin which is active against vaccinia and HSV, or 4-fluoroimidazole, 4-fluoroimidazole-5-carboxylic acid, 4-fluoroimidazole-5-carboxamide, 5-

fluoro-1-B-D-ribofurano- sylimidazole-4-carboxamide, 5-amino-1-B-D-ribofuransyl-
imidazole-4-carboxamide, poly (I).multidot.poly (C), sinefungin, iododeoxyuridine, 9-(2-
hydroxy-ethoxymethyl) guanine, gliotoxin, distamycin A, netropsin, congocidine, cordycepin, 1-
B-D-arabinofuransylthymine, 5,6-di-hydroxy-5-azathymidine, pyrazofurin, toyocamycin, or
5 tunicamycin,
an inhibitor of fungal chitin synthetase such as polyoxin D, nikko-mycin Z, or
nikkomycin X,
an impermeant antifungal agent such as ezomycin A.sub.1, A.sub.2, B.sub.1, B.sub.2,
C.sub.1, C.sub.2, D.sub.1, or D.sub.2 or platenocidin, septacidin, sinefungin, A9145A, A9145C,
10 or thraustomycin,
an inhibitor of central nervous system carbonic anhydrase such as methazolamide, or 2-
benzoylimino-3-methyl-.DELTA..sup.4 -1,3,4-thiadiazoline-5-sulfonamide substituted at the
benzoyl group with 3,4,5-trimethoxy, 2,4,6-trimethoxy, 2,4,5-trimethoxy, 4-chloro, 4-bromo, 4-
iodo, or hydrogen,
15 an inhibitor of dopamine-B-hydroxylase during the synthesis of norepinephrine and
epinephrine such as fuscic acid, 5-(3',4'-dibromobutyl)picolinic acid, 5-(3'-bromobutyl)
picolinic acid, 5-(3',4'-dichlorobutyl)picolinic acid, YP-279, benxyloxyamine, p-
hydroxybenzyloxyamine, U-21,179, U-7231, U-6324, U-0228, U-5227, U-10,631, U-10,157, U-
1238, U-19,963, U-19,461, U-6628, U-20,757, U-19,440, U-15,957, U-7130, U-14,624, U-
20 22,996, U-15,030, U-19,571, U-18,305, U-17,086, U-7726, dimethyldithiocarbamate,
diethyldithiocarbamate, ethyldithiocarbamate, 2-mercaptoethylguanidine, thiophenol, 2-
mercaptoethylamine, 3-mercaptoethylguanidine, 3-mercap- toprbpyl-N-methylguanidine, 2-
mercaptoethanol, 2-mercaptoethyl-N-methylguanidine, 2-mercaptoethyl-N,N'-
dimethylguanidine, 4,4,6-trimethyl-3,4-dihydropyrimidine-2-thiol, N-phenyl-N'-3-(4H-1,2,4-
25 trizolyl)thiourea, methylspinazarin, 6,7-dimethylspinazarin, 7-O-methy-spinochrome B, 6-(3-
hydroxybutyl)-7-O-methylspinachrome B, aquayamycin, chrothiomycin, frenoclicin, N-n-butyl-
N'-3-(4H-1,2,4-triazolyl) thiourea, propylthiouracil, mimosine, mimosinamine, or mimosinic acid,
an inhibitor of histidine decarboxylation during the synthesis of histamine such as .sup.2 -
hydroxy-5-carbomethoxybenzyloxyamine, 4-toluene-sulfonic acid hydrazide, 3-hydroxy
30 benzyloxyamine, hydroxylamine, aminooxyacetic acid, 4bromo-3-hydroxybenzyloxyamine
(NSD-1055), rhodanine substituted in the 3 position with p-chlorophenethyl, p-chlorobenzyl, p-
methylthiobenzyl, p-methylbenzyl, p-fluorobenzyl, amino, 3,4-dichlorobenzyl, p-bromobenzyl,
p-methoxybenzyl, p-bromoanilino, p-iodoanilino, p-chloroanilino, p-toluidino, anilino, 2,5-
dichloroanilino, dimethylamino, or p-methoxyphenyl; 2-mercaptobenzimidazole-1,3-dimethylol,
35 4-bromo-3-hydroxy -benzoic acid, 4-bromo-3-hydroxybenzyl alcohol, 4-bromo-3-hydroxy-
hippuric acid, (R,S)-.alpha.-fluoromethyl- histidine, (S)-.alpha.-fluoromethylester, L-histidine
ethyl ester, L-histidinamide, D,L-3-amino-4-(4-imidazolyl)-2-butanone, 2-bromo-3-
hydroxybenzyloxyamine, 5-bromo-3-hydroxybenzyloxyamine, 4,6-dibromo-3-
hydroxybenzyloxyamine, aminooxypropionic acid, benzyloxyamine, 4-bromo-3-

benzenesulfonyloxybenzyloxyamine, 3',5,7-trihydroxy-4',6-dimethoxyisoflavone, lecanoric acid, N-(2,4-dihydroxybenzoyl)-4-aminosalicylic acid, or 3',5,7-trihydroxy-4',8-dimethoxyisoflavone,

an pharmaceutical agent of drug that appear in Physicians Desk Reference, Edward R. Barnhart, 41th ed., 1987, Medical Economics Company Inc., N.J.; USAN and the Dictionary of Drug Names, ed. by Mary C. Griffiths, The United States Pharmacopiedial Convention, (1986); and The Pharmacological Basis of Therapeutics, ed. by A.G. Gilman, L. Goodman, A. Gilman, 7th ed., (1985), MacMillan Publishing Co., N.Y., N.Y.;

a centrally acting converting enzyme inhibitor such as captopril,

an antibacterial agent such as penicillin, cephalosporin, or cephamycin, with B-lactamase resistance,

an agent which blocks bacterial synthesis of tetrahydrofolate such as a sulfonamide (an analogue of p-aminobenzoic acid) including sulfanilamide, sulfadiazine, sulfamethoxazole, sulfisoxazole, or sulfacetamide

an inhibitor of dihydrofolate reductase including pyrimethamine, cycloguanil, trimethoprin, isoaminopterin, 9-oxofolic acid, or isofolic acid,

a bactericidal agent such as nalidixic acid or oxolinic acid,

an inhibitor of bacterial protein synthesis such as vancomycin, an aminoglycoside, erythromycin, tetracyclin, or chloramphenicol,

an inhibitor of viral DNA polymerase such as vidarabine,

tuberculostatic or tuberculocidal agent such as isoniazid or aminosalicylic acid,

an anthelmintic agent such as oxamniquine, piperazine, metronidazole, diethylcarbamazine, paromomycin, niclosamide, bithionol, metrifonate, hycanthone, dichlorophen, or niclosamide,

an H₂-blocking agent such as cimetidine or ranitidine,

an agent which blocks release of norepinephrine such as sotalol, guanethidine, pindolol, pronethalol, KO 592, practolol, oxprenolol, or pronethalol,

a xanthine oxidase inhibitor such as allopurinol, thioinosinate, 5,7-dihydroxypyrazolo [1,5-a] pyrimidine substituted at the 3 position with hydrogen, nitro, bromo, chloro, phenyl, 3-pyridyl, p-bromophenyl, p-chlorophenyl, p-acetylanilino, p-tolulyl, m-tolulyl, naphthyl, or 3,4-methylenedioxyphenyl;

8-(m-bromoacetamidobenzylthio)hypoxanthine, 8-(m-bromoacetamidobenzylthio)hypoxanthine, guanine substituted at the 9 position with phenyl, 4-chlorophenyl, 3-chlorophenyl, 3,4-dichlorophenyl, 4-methoxyphenyl, 3,4-dimethoxyphenyl, 4-dimethylaminophenyl, 4-aminophenyl, 3-aminophenyl, 3-trifluormethylphenyl, 4-benzamido, 4-carboxylphenyl, 4-methylphenyl, 4-ethylphenyl, 3-methylphenyl, B-naphthyl, or 4-ethoxyphenyl;

4,6-dihydroxypyrazolo [3,4-d] pyrimidine, 4-trifluoromethylimidazoles substituted at the 2 position with phenyl, p-chlorophenyl, p-methoxyphenyl, p-acetylanilino, p-nitrophenyl, p-dimethylaminophenyl, p-cyanophenyl, p-fluorophenyl, p-carboxyphenyl, m-chlorophenyl, 3,4-dichlorophenyl, 4-pyridyl, 3-pyridyl, 2-quinolyl, 6-quinolyl, 4-quinolyl, 7-quinolyl, 2-pyrazinyl, or 1-(2-pyridyl-4-trifluoromethyl-5-bromoimidazolyl); 5-(4-pyridyl)-1,2,4-triazoles substituted at

the 5 position with 4-pyridyl, 3-pyridyl, 2-pyridyl, phenyl, p-chlorophenyl, m-chlorophenyl, p-sulfonamidophenyl, 3,5-dichlorophenyl, 3,5-dicarboxyphenyl, 6-quinolyl, 2-furyl, 4-pyridazinyl, 2-thienyl, 2-pyrimidinyl, 4-pyrimidinyl, or 4-pyrazinyl; difunisal, 4(or 5)-(2-aminoethylthio-azo)imidazole-5(or 4)-carboxamide, 4 (or 5)-diazimidazole-5(or 4)-carboxamide, or S-5(or 4)-carbamoyl-4(or 5)-imidazolyl azo! cysteine,

an agent which inhibits DNA synthesis such as a bis-thiosemicarbazone, 3,5-diisopropylsalicyl- hydroxamic acid, 4-hydroxybenzoylhydroxamic acid, 3-methylsalicylhydroxamic acid 2,5-dihydroxybenzoylhydroxamic acid, or 2-hydroxy-3,4,5-trimethoxybenzoylhydroxamic acid; or which inhibits nucleotide synthesis such as N-(phosphoacetyl)-L-aspartate which inhibits asparatate transcarbamylase during pyrimidine synthesis, or azaserine or 6-diazo-5-oxo-L-norleucine which inhibits purine synthesis at the phosphoribosyl-formyl-glycineamidase synthetase step; or which is an antifolate such as methotrexate, 2,4-diamino-5-benzyl-6-(4-phenylbutyl) pyrimidine, 2,4-diamino-5-phenyl-6-(4-phenylbutyl) pyrimidine, 2,4-diamino-5-phenyl-6-(3-anilinopropyl) pyrimidine, 2-amino-4-hydroxy-5-phenyl-6-(3-p-aminobenzoylglutamic acid propyl) pyrimidine, N-(p-oo(2,4-diamino-6-quinazoliny)l)methyl-methylamino- benzoyl-L-glutamic acid, N-5p-5,2,4-diamino-5-methylquinazoliny)l)methylamino!benzoyl-L-aspartic acid, N-5p-5,2(2-amino-4-hydroxy-6-quinazoliny)l) methyl-!methylamino! benzoyl!-L-glutamic acid, 2,4-diaminoquinazolines: CCNSC 105952, CCNSC 112846, CCNSC 121346, CCNSC 122761, CCNSC 122870, CCNSC 529859, CCNSC 529860, or CCNSC 529861; 8-aza GMP, 7-deaza-8-aza GMP, 2'-dGMP, B-D-arabinosyl GMP, pentopyranine A-G, B-ribofuranosyl-1,3-oxazine-2,4-dione, pyrazofurin, 6-(p-chloroacetyl-anilinomethyl)-5-cetylvinylanilinomethyl)-5-(p-chlorophenyl)-2,4-diaminopyridine, 6-(p-chloroacetyl- ethylanilino-methyl)-5-(p-chlorophenyl)-2,4-diamino pyridine, 6-(p-chlorophenylbutylanilinomethyl)-5-(p-chlorophenyl)-2,4-diamino pyridine, p-(2,6-diamino-1,2-dihydro-2, 2-dimethyl- S-triazin-1-yl) phenylpropionyl sulfanilylfluoride or variants of the propionamide bridge of acrylamido, N-ethylsulfonamido, N-ethylcarboxamido, oxyacetamido, or oxythyloxy; or which inhibits purine or pyrimidine synthesis such as xylosyladenine, 6-azauridine, 5-aminouridine, 5-azauric acid; or which inhibits nucleotide interconversion such as hadacidin, 6-mercaptapurine, azathioprine, nitro-dUMP, psicofuranine, decoyinine, 5-fluorouracil, 5-fluorodeoxyuridine, shadowmycin; or which inhibits nucleotide utilization such as cytosine arabinoside, arabinosyladenine; or which becomes incorporated into polynucleotides such as 8-azaguanine, tubercidine, toyocamycin, sangivamycin, formycin, 7-deazainosine, 8-azainosine, or 7-thia-7, 9-dideazainosine; or which is a glyoxalase inhibitor such as Glyo-I, or Glyo-II,

an agent which blocks synthesis of prostaglandin A₂ which effects platelet aggregation such as salicylic acid, pyrogallol, 5,8,11,14-eicosatetraenoic acid, .alpha.-naphthol, guaiacol, propylgallate, nordihydroguaiaretic acid, N-0164, benzydamine, 9,11-azoprostanoic acid, 2-isopropyl-3-nicotinylindole,

an agent which blocks prostaglandin synthetase such as indomethacin, sulindac, tolmetin,

- mefenamic acid, ibuprofen, naprozen, fenoprofen, fluribiprofen, ketoprofen, meclofenamic acid, flufenamic acid, niflumic acid, benzydamine, oxyphenbutazone, aspirin, acetaminophen, salicylamide, O-carboxydiphenylamine, tolectin, diclofenac, 2,7-dihydroxynaphthalene, 5-(4-chlorobenzoyl)-1-methylpyrrole-2-acetic acid, 5-(4-methylbenzoyl)-1,4-dimethylpyrrole-2-acetic acid, 5-(4-chlorobenzoyl)-1,4-dimethylpyrrole-2-acetic acid, 5-(4-fluorobenzoyl)-1,4-dimethylpyrrole-2-acetic acid, 5-(4-chlorobenzoyl)-1,4-dimethylpyrrole-2-(2-propionic acid), 5,6-dehydroarachidonate, 11,12-dehydroarachidonate, or 5,8,11,14-eicosatetraynoate; or of an agent which blocks lipoxygenase or blocks leukotriene action such as BW755C, FPL 55712, or U-60,257
- 10 an antiarrhythmic agent such as procainamide or quinidine,
an inhibitor of hepatic synthesis of Vitamin K dependent clotting factors such as warfarin sodium, dicumarol, 4-hydroxycoumarin, phenprocoumon, or acenocoumarol,
an agent which relaxes vascular smooth muscle such as hydralazine, minoxidil, or isoxsuprine,
- 15 a Na.sup.+ -K.sup.+ -ATPase inhibitor such as digitoxigenin, digoxigenin, cymarol, periplogenin, or strophanthidiol, or ouabain glycosides, cardenolides, or basic esters, or ICI-63,632, ICI-63,605, ICI-62-655, ICI-62,838, ICI-69,654, ICI-58,622, ICI-61,374, ICI-57,267, ICI-61,424, ICI-61,411, ICI-65,199, ICI-70,898, ICI-70,899, ICI-70,900, ICI-70,901, ICI-62,966, ICI-65,210, ICI-63,116, ICI-62,936, ICI-65,551, ICI-63,978, ICI-62,276, ICI-63,056, ICI-67,135,
- 20 ICI-67,167, ICI-67,134, ICI-67,875, ICI-67,880, or ICI-61,558,
a calcium channel blocker such as prenylamine, verapamil, fendiline, gallopamil, cinnarizine, tiapamil, diltiazem, bencyclan, or nifedipine; or an agent which stabilizes calcium binding to cellular calcium stores and thereby inhibits the release of this calcium by contractile stimuli such as 8-(N,N-diethylamino)-octyl 3,4,5-trimethoxybenzoate (TMB-8),
- 25 a monoamine oxidase inhibitor such as tranlycypromine, phenylethylamine, trans-cinnamic acid, phenelzine, or isocarboxazid,
a benzodiazepine compound such as clorazepate,
valproic acid,
an agent which causes repression of the synthesis of HMG-CoA reductase such as 20-
- 30 .alpha.-hydroxycholesterol, 22-ketocholesterol, 22-.alpha.-hydroxycholesterol, 25-hydroxycholesterol, 22-B-hydroxycholesterol, 7-.alpha.-hydroxycholesterol, 7-B-hydroxycholesterol, 7-ketocholesterol, or kryptogenin; or of an agent which inhibits HMG-CoA reductase such as, lorelco; or of an agent which inhibits lipolysis such as 5-methylpyrazole -3-carboxylic acid (U-19425), nicotinic acid, uridine, inosine, 3,5-dimethylisoxazole (U-21221),
- 35 3,5-dimethylpyrazole, prostaglandin E.sub.2, eritadenine, or eritadenine isoamyl ester; or of an agent which inhibits lipogenesis such as ascofuranone, (-)-hydroxycitrate, or tetrol-yl-CoA; or of an agent which is hypocholesterolemic such as lentysine; or of an agent which lowers triglycerides such as lopid; or of an agent which is an inhibitor of acetyl-CoA carboxylase during lipogenesis such as 2-methyl -2- δ p-(1,2,3,4-tetrahydro-1-naphthyl)-phenoxy!-propionate

(SU13437), .sup.2 -(p-chlorophenoxy)-2-methylpropionate, kynurenate, xanthurenate, kynurenine, 3-hydroxyanthranilate, or 2-methyl-2-δp-(p-chlorophenyl)phenoxy! propionate; or of an agent which is an inhibitor of hepatic B-lipoprotein production such as orotic acid,

a vasodilator such as WS-1228A, or WS-1228B; or of an anti-inflammatory agent such as amicomacin A,

a protease inhibitor such as leupeptin; or which is an inhibitor of pepsin such as a pepstatin, a pepstanone, or a hydroxyepstatin,

an inhibitor of cell surface enzymes such as bestatin, amastatin, forphenicine, ebelactone, or forphenicin,

a phosphodiesterase inhibitor such as theophyllineacetic acid, theophylline, dyphylline, disodium cromoglycate, 6-n-butyl-2,8-dicarboxy-4,10-dioxo-1,4,7,10-tetrahydro-1,7-phenanthrolin, 2-chloroadenosine, dipyridamole, EG 626, AY-17,605, AY-17,611, AY-22,252, AY-22,241, cis-hinokiresinol, oxy-cis-hinokiresinol, tetrahydro-cis- hinokiresinol, trans-hinokiresinol, dehydrodicaffeic acid, 2,6,4'-trihydroxy-4-methoxybenzophenone, p-hydroxyphenyl crotonic acid, papaverine, 3-(5-tetrazolyl)-thioxanthone-10,10-dioxide, 3-carboxythioxanthone-10,10-dioxide, W-7, HA-558, MY-5445, OPC-3689, OPC-13135, or OPC-13013, reticulol, PDE-I, or PDE-II,

an inhibitor of tyrosine hydroxylase, the enzyme catalyzing the rate-limiting reaction in the biosynthesis of norepinephrine, such as azadopamine, isopropylazadopamine, dimethylazadopamine; triphenolic compounds such as n-propylgallate; diphenolic benzoic acid derivatives such as 3,4-dihydroxybenzoic acid; phenylcarbonyl derivatives such as 3,4-dihydroxybenzaldehyde, arterenone, or adrenalone H 22/54, 3-iodo-L-tyrosine, D,L-.alpha.-methyl-p-tyrosine, L-3-iodo-.alpha.-methyltyrosine, 3-bromo-.alpha.-methyltyrosine, gentistic acid, 3-chloro-.alpha.-methyltyrosine, phenylalanine derivatives, 3,5-diiodo- L-tyrosine, 3,5-dibromo-L-tyrosine, 3-bromo-.alpha.-methyl-L- tyrosine, 3-fluro-.alpha.-methyl-L-tyrosine, catechol analogues, 3,4-dihydroxyphenylethylacetamide, 3,4-dihydroxyphenylisopropylacetamide, 3,4-dihydroxyphenylbutylacetamide, 3,4-di-hydroxyphenylisobutylacetamide, D,L-.alpha.-methylphenylalanine, D,L-3-iodophenylalanine, D,L-4-iodophenylalanine, D,L-.alpha.-methyl-3-iodophenylalanine, D,L-a-methyl-3-bromophenylalanine, D,L-.alpha.-methyl-3-chlorophenylalanine, D,L-.alpha.-methyl-3-fluorophenylalanine, mimosine, mimosinamine, mimosinic acid, 7-O-methylspinochrome B, 6-(3-hydroxybutyl)-7-O-methylspinachrome B, aquayamycin, chrothiomycin, frenolicin, fuscic acid, pentylpicolinic acid, dopstatin, methylspinazarin, 6,7-dihydroxymethylspinazarin, 3-ethyl-.alpha.-methyltyrosine, 3-methyl-.alpha.-methyltyrosine, 3-isopropyl-x-methyltyrosine, 3-allyl-.alpha.-methyltyrosine, 3-δ4-hydroxy-3-(2-methylallyl)-phenyl!-2-methylalanine,, 3-δ3-(2,3-epoxypropyl)-4-hydroxyphenyl!-2-methylalanine, 3-isobutyl-.alpha.-methyltyrosine, 3-methylvinyl-.alpha.-methyltyrosine, 5-methyl-6,7-diphenyltetrahydropterin, 3-(2,3-dihydro-2,2-dimethyl-5-benzofuranyl!-2-methylalanine, 3-δ2,3-dihydro-2,2-dimethyl-5-benzofuranyl!-2-methylalanine, .alpha.-methyl-dopa, or ethyl-3-amino-4H-pyrrolo δ3,4c! isoxazole carboxylate, and

proteins including enzymes and hormones such as insulin, erythropoietin, interleukin 2, interferon, growth hormone, atrial natriuretic factor, tissue plasminogen activator.

37. The method of synthesis of the compound of claim 1 wherein the C moiety comprises at least one of the group of herbicides, fungicides, miticides, nematocides, fumigants, growth regulators, repellants, defoliants, rodenticides, molluscicides, algicides, desiccants, antehelminthics, and bactericides.

38. The method of synthesis of the compound of claim 37 wherein the C moiety is one from the those given in Chemical Week Pesticides Register, Robert P. Ovellette and John A. King, 1977, McGraw-Hill Book Company.

39. A method of synthesis of a chemical compound having the formula $(A-B-C)_x-P-E_y$ where the A is a chemiluminescent moiety,

B is an energy acceptor moiety, and

C is a biologically active moiety, and

P is a substrate

E is an enzyme and x and y are integers

comprising the steps of

forming a benzophenone,

forming a diaryl ethylene,

attaching a phthalimide moiety to at least one of the aryl groups of the ethylene to form a phthalimide-ethylene conjugate,

condensing two ethylene-phthalimide conjugates to form a phthalimide-pentadiene conjugate,

converting the phthalimide to the phthalhydrazide by reaction with hydrazine to form a carrier compound, and

reacting the carrier compound with a biologically active moiety to form a corresponding conjugate,

reacting $A-B-C$ with a polymer to form $(A-B-C)_x-P$, and

reacting E with $(A-B-C)_x-P$ to form $(A-B-C)_x-P-E_y$.

40. The method of synthesis of the compound of claim 39 wherein the compound provides controlled extra cellular release of the C moiety.

41. The method of synthesis of the compound of claim 39 wherein the C moiety comprises at least one of drugs and proteins including enzymes and hormones.

42. The method of synthesis of the compound of claim 41 wherein the C moiety comprises at

least one insulin, erythropoietin, interleukin 2, interferon, growth hormone, atrial natriuretic factor, tissue plasminogen activator, an anti-inflammatory drug, an antihypertensive drug, an inotropic drug, and a contraceptive drug.

5 43. The method of synthesis of the compound of claim 40 wherein extraacellular drug release occurs when the prodrug reacts with cellular free radicals via a mechanism involving chemiluminescence, photochromism, and intramolecular energy transfer.

10 44. The method of synthesis of the compound of claim 41 wherein the pharmaceutical agent is at least one of the group of antilipidemic drugs, anticholesterol drugs, contraceptive agents, anticoagulants, anti-inflammatory agents, immuno-suppressive drugs, antiarrhythmic agents, antineoplastic drugs, antihypertensive drugs, epinephrine blocking agents, cardiac inotropic drugs, antidepressant drugs, diuretics, antifungal agents, antibacterial drugs, anxiolytic agents, sedatives, muscle relaxants, anticonvulsants, agents for the treatment of ulcer disease, agents for
15 the treatment of asthma and hypersensitivity reactions, antithrombotic agents, agents for the treatment of muscular dystrophy, agents to effect a therapeutic abortion, agents for the treatment of anemia, agents to improve allograft survival, agents for the treatment of disorders of purine metabolism, agents for the treatment of ischemic heart disease, agents for the treatment of opiate withdrawal, agents which activate the effects of secondary messenger inositol triphosphate,
20 agents to block spinal reflexes, and antiviral agents including a drug for the treatment of AIDS.

25 45. The method of synthesis of the compound of claim 43 wherein the C moiety is released by an oxidation reduction reaction with the target cell's electron carriers or by reaction with free radicals produced as a consequence of electron transport.

46. The method of synthesis of the compound of claim 43 wherein A represents a functionality which undergoes at least one of

an oxidation reduction reaction where electrons are transferred directly between A and the target cell's electron carriers, and

30 a reaction with free radicals of oxygen which are produced as a consequence of electron transport

such that an excited state is produced in A as a consequence of its participation in one of these reactions.

35 47. The method of synthesis of the compound of claim 46 wherein A undergoes intramolecular energy transfer from its own excited state to the B functionality which is an energy acceptor.

48. The method of synthesis of the compound of claim 47 wherein upon receiving energy

from A, B achieves an excited state which relaxes through heterolytic cleavage of the covalent bond of B with C where C is a drug moiety which is released into the environment.

49. The method of synthesis of the compound of claim 39 wherein the chemiluminescent molecule comprises at least one of the group of

molecules undergoing reaction involving peroxides and oxygen free radicals,

molecules undergoing reaction involving oxidation or reduction, and

molecules undergoing both reaction with peroxides and oxygen free radicals followed by an oxidation or reduction reaction.


50. The method of synthesis of the compound of claim 49 wherein the chemiluminescent molecule comprises at least one of the group of luminol and its derivatives, lucigenin and its derivatives, Lophine and its derivatives, acridinium esters and acridans, tetraphenylpyrrole, phthalhydrazides, acyloins, biacridinium salts, vinylcarbonyls, vinylnitriles, tetrakis (dimethylamino) ethylene, acylperoxides, indoles, tetracarbazoles and active oxalates.

51. The method of synthesis of the compound of claim 49 wherein the chemiluminescent molecule comprises at least one of the group of ruthenium chelates 2, 6-diaminopyrene, or cation radicals and molecules which follow a Chemically Initiated Electron Exchange Luminescence mechanism such as certain dioxetans and dioxetanones.

52. The method of synthesis of the compound of claim 49 wherein the chemiluminescent molecule comprises at least one of the group of dioxene derivatives and other compounds that form a dioxetan by reaction with superoxide and then produce efficient chemiluminescence by a CIEEL mechanism.

53. The method of synthesis of the compound of claim 49 wherein the **chemiluminescent molecule** comprises at least one of the group of

2,6-diaminopyrene



Aminophthalhydrazide




TABLE 1

5 Representative Chemiluminescent Molecules

Name	Structure
------	-----------

10

TABLE 1 continued

Representative Chemiluminescent Molecules









5	Name	Structure
	Dioxene	
	Imidazole derivatives	
	Sulfonyloxamides	
	Indole derivatives	
	Tetrakis(dialkylamino)-ethylene	
	2,5,7,8-tetraoxabicyclo-[4.2.0.] octane	
	Dioxetan	
	Lucigenin	

TABLE 1 continued


Representative Chemiluminescent Molecules

5


Name

Structure


Lophine




Acridinium esters




Active oxalate




Tris-2,2'-bipyridinedi-chlororuthenium (II)



Dioxetanone



Diphenyl peroxide



10

54. The method of synthesis of the compound of claim 39 wherein the B moiety is a photochromic compound.

55. The method of synthesis of the compound of claim 54 wherein the photochromic compound comprises one which demonstrate photochromic behavior with electromagnetic radiation and bleaching agents.

5


56. The method of synthesis of the compound of claim 55 wherein the A functionality is chemiluminescent, and the B functionality is such that the photodissociative drug release spectrum of B overlaps the chemiluminescence spectrum of A.

10 57. The method of synthesis of the compound of claim 54 wherein the photochromic compound comprises a cationic dye.


58. The method of synthesis of the compound of claim 57 wherein the cationic dye comprises at least one of a di and triarylmethane dyes, triarylmethane lactones and cyclic ether dyes,
15 cationic indoles, pyronines, phthaleins, oxazines, thiazines, acridines, phenazines, and anthocyanidins, and cationic polymethine dyes and azo and diazopolymethines, styryls, cyanines, hemicyanines, dialkylaminopolyenes, and other related dyes.

20 59. The method of synthesis of the compound of claim 57 wherein the **cationic dye** comprises at least one of

2,6-diaminopyrene




Aminophthalhydrazide




25


Dioxene




Imidazole derivatives




Sulfonyloxamides




Indole derivatives




Tetrakis(dialkylamino)-ethylene




2,5,7,8-tetraoxabicyclo-[4.2.0.]octane




Dioxetan




Lucigenin




Lophine




Acridinium esters




Active oxalate




Tris-2,2'-bipyridinedi-chlororuthenium (II)



Dioxetanone



Dipheyl peroxide



5

60. The method of synthesis of the compound of claim 39 wherein the C moiety is any molecule which exhibits bleaching behavior with the B moiety and has an increased therapeutic effect or therapeutic ratio as a consequence of its delivery as part of a prodrug.


61. The method of synthesis of the compound of claim 39 wherein the C moiety has a nucleophilic group that bonds to the B moiety.

62. The method of synthesis of the compound of claim 61 wherein the C moiety is derivatized to have a nucleophilic group that bonds to the B moiety.


5 63. The method of synthesis of the compound of claim 62 wherein the C moiety is derivatized by at least one of the nucleophilic groups comprising cinnamate, sulfite, phosphate, carboxylate, thiol, amide, alkoxide, or amine.

10 64. The method of synthesis of the compound of claim 39 wherein the C moiety is at least one of the group of


• Captopril




Prostaglandin E₂




2,3-dichloro- α -methylbenzylamine




3'-deoxy-S-adenosyl-L-homocysteine




Sinefungin




3,5-diiodo-4-hydroxybenzoic acid




6,6'-dithiobis (9-B-D-ribofuranosylpurine)




γ -aminobutyric acid




Gabaculine




N-(5'-phosphopyridoxy)-
4-aminobutyric acid




4-amino-hex-5-enoic acid




Baclofen




Adenosine




3-hydroxy-3-methyl-
glutarate




Campactin




But-3-ynoyl-CoA




Suramin




L-3-iodotyrosine




L-3-iodo-α-methyltyrosine




Disodium cromoglycate




Adenosine
3',5'-cyclic monophosphate




D,L-B-(5-hydroxy-3-indolyl)- α -hydrazinopropionic acid




D,L- α -hydrazino- α -methyl dopa




α -methyl dopa




5-(3,4-dihydroxycinnamoyl)salicylic acid




N-(phosphonacetyl)-L-aspartate




P-glycolohydroxamate




5-(p-sulfamylphenyl)-azosalicylic acid




Coformycin



Formycin B




Thioinosinate




Phosphonoformate




Phosphonoacetate




Ridavirin




Sotalol



Cimetidine




Fuscaric acid




2-mercaptoethylamine




Mimosine




U-7130




Iproniazid




Trans-4-aminocrotonic acid




NSD 1055




Nicotinic acid




Kynurenic acid




Lentysine




Orotic acid




Polyoxin D



Cephalosporin



Penicillin



5

65. The method of synthesis of the compound of claim 39 wherein the A-B-C moieties are attached to P by a bond between P and at least one of A and B.

10

66. The method of synthesis of the compound of claim 39 wherein the E moieties are attached to (A-B-C)_x-P by a bond between E and at least one of A, B, and P.

67. The method of synthesis of the compound of claim 39 wherein the E moieties are enzymes that react with a desired substrate and form substances that cause the release of C from A-B-C.

15

68. The method of synthesis of the compound of claim 39 wherein the E moieties are enzymes that react with a desired substrate and form peroxide or free radicals that cause the

release of C from A-B-C.

69. The method of synthesis of the compound of claim 67 wherein the E moiety, substrate, and C moiety are at least one of the group of

glucose oxidase, glucose, and insulin, and
xanthine oxidase, xanthine, and tissue plasminogen activator (TPA).

70. A method of synthesis of a chemical compound having the formula $(A-B-C)_x-P$

where the A is a chemiluminescent moiety,

B is an energy acceptor moiety, and

C is a biologically active moiety, and

P is a substrate and x is an integer

comprising the steps of

forming a benzophenone,

forming a diaryl ethylene,

attaching a phthalimide moiety to at least one of the aryl groups of the ethylene to form a phthalimide-ethylene conjugate,

condensing two ethylene-phthalimide conjugates to form a phthalimide-pentadiene conjugate,

converting the phthalimide to the phthalhydrazide by reaction with hydrazine to form a carrier compound, and

reacting the carrier compound with a strong base such as an alkali hydride and the biologically active moiety to form a corresponding conjugate,

reacting A-B-C with a polymer to form $(A-B-C)_x-P$.

71. The method of synthesis of the compound of claim 1 wherein one or more of the moieties can be modified to further candidate components by addition of functional groups.


72. The method of synthesis of the compound of claim 71 wherein the groups comprise at

least one of alkyl, cycloalkyl, alkoxycarbonyl, cyano, carbamoyl, heterocyclic rings containing C, O, N, S, sulfo, sulfamoyl, alkoxysulfonyl, phosphono, hydroxyl, halogen, alkoxy, alkylthiol, acyloxy, aryl, alkenyl, aliphatic, acyl, carboxyl, amino, cyanoalkoxy, diazonium, carboxyalkylcarboxamido, alkenylthio, cyanoalkoxycarbonyl, carbamoylalkoxycarbonyl, alkoxy carbonylamino, cyanoalkylamino, alkoxycarbonylalkylamino, sulfoalkylamino, alkylsulfamoylalkylamino, oxido, hydroxy alkyl, carboxy alkylcarbonyloxy, cyanoalkyl, carboxyalkylthio, arylamino, heteroarylamino, alkoxycarbonyl, alkylcarbonyloxy, cyanoalkoxy, alkoxycarbonylalkoxy, carbamoylalkoxy, carbamoylalkyl carbonyloxy, sulfoalkoxy, nitro, alkoxyaryl, halogenaryl, amino aryl, alkylaminoaryl, tolyl, alkenylaryl, allylaryl, alkenyloxyaryl, allyloxyaryl, cyanoaryl, carbamoylaryl, carboxyaryl, alkoxycarbonylaryl, alkylcarbonyoxyaryl,

sulfoaryl, alkoxysulfoaryl, sulfamoylaryl, and nitroaryl.

73. The method of synthesis of the compound of claim 1 wherein the compound has the structure of general formula

5



74. The method of synthesis of the compound of claim 73 wherein the functionality A is at least one of aminophthalhydrazide derivatives, sulfonyloxamides and active oxalates,

the functionality B is at least one of 1,1,5,5-tetrakisarylpentadiene and 1,1,5-

10 trisarylpentadiene derivatives,

the functionality C is a drug molecule such as Foscarnate, or ddc;, and


R is a functional group, and

L is a linker such as an aliphatic chain between A and B.


15 75. The method of synthesis of the compound of claim 74 wherein the L functionality is between one 20 carbon atoms.

76. The method of synthesis of the compound of claim 1 wherein B is a 1,1,5-trisarylpentadiene derivative and the compound has the formula

20





77. The method of synthesis of the compound of claim 1 wherein A is a sulfonyloxamide or active oxalate and the compound has the formula




78. The method of synthesis of the compound of claim 1 wherein a luminol derivative is
5 directly attached through one or more amino groups to the aryl groups of a photochromic dye.


79. The method of synthesis of the compound of claim 78 wherein
C comprises the formula of at least one of




10 ddc , and A-B comprises the formula of at least one of




YY99811-1



6a




GZW2-33-1
 $C_{37}H_{38}N_5O_2 \cdot ClO_4$
 M.W. 684.20



GZW1-98-2
 $C_{49}H_{41}N_6O_4 \cdot BF_4$
 M.W. 864.71

, and

MTLJ-1




5

80. The method of synthesis of the compound of claim 78 wherein the compound comprises

the formula

MTLJ-1-Foscarnet



81. The method of synthesis of the compound of claim 1 wherein the hydrolyzable group that protects phthalhydrazide is at least one of acetyl and t-butyloxycarbonyl.

82. The method of synthesis of the compound of claim 1 wherein the aminophthalimide-substituted precursors for the dye are prepared through amination of an aryl halide such as palladium-catalyzed amination of aryl halides.

83. The method of synthesis of the compound of claim 1 wherein halo-substituted aryl groups of a starting B moiety or an intermediate are coupled with the aminophthalimide by methods such as the aryl amination under palladium catalysis to form the aminophthalimide-substituted precursors for the dye.

84. The method of synthesis of the compound of claim 1 wherein halo-substituted aryl groups of a starting phthalimide or an intermediate are coupled with the amino-substituted dye by methods such as the aryl amination under palladium catalysis to form the aminophthalimide-substituted precursors for the dye.

85. The method of synthesis of the compound of claims 83 and 84 wherein amino-substituted aryl groups are obtained by the amination of the halo-substituted compounds with an imine such as benzophenoneimine.

86. The method of synthesis of the compound of claim 1 wherein the aminophthalimide-attached dye is formed by the condensation of two aminophthalimide-attached ethylene molecules by reaction with triethyl orthoformate and a strong acid such as perchloric acid in acetic anhydride or acetic acid.

87. The method of synthesis of the compound of claim 1 wherein during the step of converting the phthalimide moiety to the aminophthalhydrazide to obtain A-B, the B moiety is protected from reaction with hydrazine by reacting with base such as sodium hydroxide, sodium methoxide and amines.

88. The method of synthesis of the compound of claim 87 wherein the phthalimide-B conjugate with a protected B moiety is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent in inert atmosphere and then treated with acid such as perchloric acid, tetrafluoroboric acid to regenerate a corresponding unaltered B moiety of the A-B conjugate.

89. The method of synthesis of the compound of claim 88 wherein A-B is reacted with one nucleophilic species of C to form A-B-C.


90. The method of synthesis of the compound of claim 1 wherein A-B is formed by starting with B comprising halo-substituted dyes, such as 1,5-bis(p-bromophenyl)-1,5-bis(p-dimethylaminophenyl)-pentadienium perchlorate.

91. The method of synthesis of the compound of claim 90 wherein cationic dyes are protected by reacting with base such as alkoxide and then coupled with the aminophthalimide by amination of aryl halide such as the palladium-catalyzed amination of aryl halide to obtain the alkoxide-protecting aminophthalimide-substituted dyes.


92. The method of synthesis of the compound of claim 91 wherein the aminophthalimide-B conjugate with a protected B moiety is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent to convert the amino-phthalimide moiety to the aminophthalhydrazide moiety and then treated with acid to generate A-B.

93. The method of synthesis of the compound of claim 1 wherein the B comprises a tetraarylpolymethine, the aminophthalhydrazide precursor is an aminophthalic acid diester and the conjugate to form A-B is amino-phthalimideluminol-tetraaryl-polymethine.

94. The method of synthesis of the compound of claim 1 wherein halo-substituted diarylketone are formed by at least one of direct acylation of arene with halo-substituted benzoyl halide under ferric chloride catalysis according to the following representative scheme



acylation according to the following representative scheme



5

95. The method of synthesis of the compound of claim 1 wherein a halo-substituted diarylketone is converted to the corresponding halo-substituted diarylketene such as halo-substituted 1,1-diarylethene.

10 96. The method of synthesis of the compound of claim 95 wherein the halo-substituted diarylketene is coupled with a precursor of amino-phthalhydrazide such as aminophthalimide, aminophthalic acid diester, by aryl amination such as the palladium-catalyzed amination of aryl halides to form the aminophthalimide-substituted 1,1-diarylethene.

15 97. The method of synthesis of the compound of claim 96 wherein the ethene is condensed with an orthoester such as triethylorthoformate in a nonaqueous solvent such as acetic anhydride, containing an acid catalyst such as perchloric acid, tetrafluoroboric acid, to form the aminophthalimide-substituted tetraarylpolymethine dye.

20 98. The method of synthesis of the compound of claim 97 wherein the aminophthalimide moiety is converted to the aminophthalhydrazide to obtain A-B.

99. The method of synthesis of the compound of claim 98 wherein the B moiety is a cationic dye that is first protected by reacting with an anion such as hydroxide, methoxide and amine and the phthalimide-B conjugate with a protected B moiety is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent in inert atmosphere and then treated with acid such as perchloric acid, tetrafluoroboric acid to regenerate a corresponding unaltered B moiety of the A-B conjugate.

30 100. The method of synthesis of the compound of claim 99 wherein A-B is reacted with one nucleophilic species of a C such as a drug 2',3'-dideoxycytidine, Foscarnet, acycloguanosine to




form A-B-C comprising a prodrug.

101. The method of synthesis of the compound of claim 95 wherein two halo-substituted diarylketene precursor compounds are condensed with an orthoester such as triethylorthoformate in a nonaqueous solvent such as acetic anhydride containing acid catalyst such as perchloric acid, tetrafluoroboric acid to form the halo-substituted tetraarylpolymethine dyes such as 1,5-bis(p-bromophenyl)-1,5-bis(p-dimethylaminophenyl)-pentadienium perchlorate.


102. The method of synthesis of the compound of claim 101 wherein the B moiety is a cationic dye that is protected by reacting with an anion such as alkoxide and then coupled with the aminophthalimide by amination of aryl halide such as the palladium-catalyzed amination of aryl halide to obtain the alkoxide-protected aminophthalimide-substituted tetraarylpolymethine dye.

103. The method of synthesis of the compound of claim 103 wherein the alkoxide-protected aminophthalimide-substituted tetraarylpolymethine dye is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent to convert the amino-phthalimide moiety to the aminophthalhydrazide moiety and then treated with acid to generate A-B comprising a luminol-tetraarylpolymethine compound.




104. The method of synthesis of the compound of claim 1 comprising the general steps given by following representative formula




4-(*N*-ethylamino)-*N*-methylphthalimide
 $\xrightarrow{\text{Pd}(\text{OAc})_2, \text{P}(t\text{-Bu})_3, t\text{-BuONa}}$




3a: $\text{R} = \text{N}(\text{CH}_3)_2$
3b: $\text{R} = \text{H}$
3c: $\text{R} = \text{OCH}_3$
3d: $\text{R} = \text{O}(\text{CH}_2)_3\text{CH}_3$
3e: $\text{R} = (\text{CH}_2)_3\text{CH}_3$







- 3a:** R = N(CH₃)₂
3b: R = H
3c: R = OCH₃
3d: R = O(CH₂)₃CH₃
3e: R = (CH₂)₃CH₃



- 4a:** R = N(CH₃)₂, X = ClO₄
4b: R = H, X = BF₄
4c: R = OCH₃, X = BF₄
4d: R = O(CH₂)₃CH₃, X = BF₄
4e: R = (CH₂)₃CH₃, X = BF₄



- 5a:** R = N(CH₃)₂, X = ClO₄
5b: R = H, X = BF₄
5c: R = OCH₃, X = BF₄
5d: R = O(CH₂)₃CH₃, X = BF₄
5e: R = (CH₂)₃CH₃, X = BF₄



105. The method of synthesis of the compound of claim 1 wherein the A functionality comprises a phthalhydrazide such as a luminol derivative and the B functionality comprises a photochromic dye wherein A is attached to aryl groups of B comprising the steps of

- 5 forming a diaryl ketone,
- forming a diaryl ketene from the diaryl ketone,
- forming a protected aminophthalhydrazide such as aminophthalimide or aminophthalic

acid diester,

adding a hydrocarbon linker to the protected aminophthalhydrazide, and

attaching the protected aminophthalhydrazide through the molecular linker to the aryl groups of diarylketene to form the precursor aminophthalimide-linked diarylketene, and reacting according to at least one of

(a) forming the A functionality from the precursor, and condensing two molecules of B precursor linked to A to form A-B, and

(b) condensing two precursor aminophthalimide-linked diarylketene molecules to form A precursor linked to B, and

forming the A functionality from the A precursor to form A-B.

106. The method of synthesis of the compound of claim 105 wherein the diaryl ketone is formed by a classical Friedel-Crafts acylation between a benzoyl halide and aryl compound with a hydrocarbon linker having a leaving group.

107. The method of synthesis of the compound of claim 106 wherein the aryl compound with a hydrocarbon linker having a leaving group comprises at least one of a halogenated-alkyl-aryl ether and a halogenated-alkyl-aryl amine wherein the halogen is the leaving group.

108. The method of synthesis of the compound of claim 107 wherein the halogenated-alkyl-aryl ether comprises 2-bromoethoxybenzene to give an aryl ketone such as 4-(2-bromoethoxy)benzophenone.

109. The method of synthesis of the compound of claim 107 wherein the halogenated-alkyl-aryl amine comprises 2-bromoethyl aminobenzene to give an aryl ketone such as 4-(2-bromoethyl amino)benzophenone.

110. The method of synthesis of the compound of claim 105 wherein the diaryl ketone is converted to the corresponding diarylketene by reacting with a methylating reagent such as a methyl Grignard reagent, methyl lithium reagent, lithium dimethylcopper reagent and then dehydration with acid.

111. The method of synthesis of the compound of claim 110 wherein the diaryl ketone is converted to the corresponding diarylketene by reacting with methylmagnesium bromide and then dehydration with acid.

112. The method of synthesis of the compound of claim 105 wherein the diaryl ketone is converted to the corresponding diarylketene by a Wittig reaction.

113. The method of synthesis of the compound of claim 105 wherein a linker is attached to the protected aminophthalhydrazide by a reaction of a nucleophilic group of the linker or protected aminophthalhydrazide with a leaving group of the linker or protected aminophthalhydrazide.

114. The method of synthesis of the compound of claim 105 wherein a linker is attached to the protected aminophthalhydrazide by reaction to form a bond between at least one of a nitrogen, oxygen, or carbon atom of the linker and at least one of a nitrogen, oxygen, or carbon atom of group of the protected aminophthalhydrazide by an addition or a substitution reaction of a leaving group.

115. The method of synthesis of the compound of claim 114 wherein a linker is attached to the protected aminophthalhydrazide by a substitution reaction of at least one of a halogen, tosylate group, ester group with a nitrogen, oxygen, or carbon atom.

116. The method of synthesis of the compound of claim 105 wherein attaching the protected aminophthalhydrazide through the molecular linker to one of the aryl groups of diarylketene to form the precursor aminophthalimide-linked diarylketene is by a reaction of a nucleophilic group of the linker or aryl group of diarylketene with a leaving group of the linker or aryl group of diarylketene.

117. The method of synthesis of the compound of claim 116 wherein a linker is attached to the aryl group of diarylketene by reaction to form a bond between at least one of a nitrogen, oxygen, or carbon atom of the linker and at least one of a nitrogen, oxygen, or carbon atom of group of the protected aminophthalhydrazide by an addition or a substitution reaction of a leaving group.


118. The method of synthesis of the compound of claim 117 wherein a linker is attached to the aryl group of diarylketene by a substitution reaction of at least one of a halogen, tosylate group, ester group with a nitrogen, oxygen, or carbon atom.

119. The method of synthesis of the compound of claim 105 wherein the precursor aminophthalimide-linked diarylketene is further reacted by condensation of two aminophthalimide-linked diarylketenes with an orthoester to form B linked to the A precursor.

120. The method of synthesis of the compound of claim 119 wherein condensing reagent is triethylorthoformate.

121. The method of synthesis of the compound of claim 119 wherein the precursor

aminophthalimide-linked diarylketene comprises at least one of the formula




12a: $R = OCH_3$

12b: $R = O(CH_2)_3CH_3$

12c: $R = (CH_2)_3CH_3$

12d: $R = N(CH_3)_2$

and the precursor of A-B comprises at least one of the formula



19a: $R = OCH_3$

19b: $R = O(CH_2)_3CH_3$

19c: $R = (CH_2)_3CH_3$

19d: $R = N(CH_3)_2$

122. The method of synthesis of the compound of claim 119 wherein the phthalimide moiety of the A precursor is converted to the phthalhydrazide A functionality by treating with hydrazine, forming A-B.


123. The method of synthesis of the compound of claims 122 wherein the B functionality is protected by reacting with an anion such as hydroxide, methoxide and amine, the A-B precursor is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent in inert atmosphere and then treated with acid such as perchloric acid, tetrafluoroboric acid to form A-B.

124. The method of synthesis of the compound of claim 105 wherein the phthalimide moiety of the A precursor of the precursor aminophthalimide-linked diarylketene is converted to the phthalhydrazide A functionality by treating with hydrazine, forming A attached to a B precursor.

125. The method of synthesis of the compound of claim 124 wherein the A-linked diarylketene is further reacted by condensation of two A-linked diarylketenes with an orthoester to form A-B.

126. The method of synthesis of the compound of claim 125 wherein condensing reagent is triethylorthoformate.

127. The method of synthesis of the compound of claim 126 wherein the A-linked diarylketene comprises at least one of the formula




18a: R=OCH₃

18b: R=O(CH₂)₃CH₃

18c: R=(CH₂)₃CH₃

18d: R=N(CH₃)₂

5 and A-B comprises at least one of the formula



20a: R=OCH₃




20b: R=O(CH₂)₃CH₃


20c: R=(CH₂)₃CH₃

20d: R=N(CH₃)₂





128. The method of synthesis of the compound of claims 123 and 125 further comprising the step of reacting the B functionality with one nucleophilic species of a C functionality such as Foscarnet to form A-B-C.


129. The method of synthesis of the compound of claim 105 comprising the general steps given by following representative formula





130. The method of synthesis of the compound of claim 105 comprising the general steps given by following representative formula





131. The method of synthesis of the compound of claim 1 wherein the A functionality comprises a phthalhydrazide such as a luminol derivative and the B functionality comprises a triarylpolymethine photochromic dye wherein A is attached to aryl groups of B comprising the steps of

- forming a diaryl ketone,
- forming a diaryl ketene from the diaryl ketone,
- forming a protected aminophthalhydrazide such as aminophthalimide or aminophthalic

acid diester,

adding a hydrocarbon linker to the protected aminophthalhydrazide, and

attaching the protected aminophthalhydrazide through the molecular linker to the aryl groups of diarylketene to form the precursor aminophthalimide-linked diarylketene, and reacting
5 according to at least one of

(a) forming the A functionality from the precursor, and condensing the A-linked diarylketene with an aryl alkene aldehyde to form A-B, and

(b) condensing the precursor aminophthalimide-linked diarylketene with an aryl alkene aldehyde to form A precursor linked to B, and

10 forming the A functionality from the A precursor to form A-B.

132. The method of synthesis of the compound of claim 131 wherein the diaryl ketone is formed by a classical Friedel-Crafts acylation between a benzoyl halide and aryl compound with a hydrocarbon linker having a leaving group.

15

133. The method of synthesis of the compound of claim 132 wherein the aryl compound with a hydrocarbon linker having a leaving group comprises at least one of a halogenated-alkyl-aryl ether and a halogenated-alkyl-aryl amine wherein the halogen is the leaving group.

20 134. The method of synthesis of the compound of claim 133 wherein the halogenated-alkyl-aryl ether comprises 2-bromoethoxybenzene to give an aryl ketone such as 4-(2-bromoethoxy)benzophenone.

25 135. The method of synthesis of the compound of claim 133 wherein the halogenated-alkyl-aryl amine comprises 2-bromoethyl aminobenzene to give an aryl ketone such as 4-(2-bromoethyl amino)benzophenone.

30 136. The method of synthesis of the compound of claim 131 wherein the diaryl ketone is converted to the corresponding diarylketene by reacting with a methylating reagent such as a methyl Grignard reagent, methyl lithium reagent, lithium dimethylcopper reagent and then dehydration with acid.

35 137. The method of synthesis of the compound of claim 136 wherein the diaryl ketone is converted to the corresponding diarylketene by reacting with methylmagnesium bromide and then dehydration with acid.

138. The method of synthesis of the compound of claim 131 wherein the diaryl ketone is converted to the corresponding diarylketene by a Wittig reaction.

139. The method of synthesis of the compound of claim 131 wherein a linker is attached to the protected aminophthalhydrazide by a reaction of a nucleophilic group of the linker or protected aminophthalhydrazide with a leaving group of the linker or protected aminophthalhydrazide.

5

140. The method of synthesis of the compound of claim 131 wherein a linker is attached to the protected aminophthalhydrazide by reaction to form a bond between at least one of a nitrogen, oxygen, or carbon atom of the linker and at least one of a nitrogen, oxygen, or carbon atom of group of the protected aminophthalhydrazide by an addition or a substitution reaction of a leaving group.

10

141. The method of synthesis of the compound of claim 140 wherein a linker is attached to the protected aminophthalhydrazide by a substitution reaction of at least one of a halogen, tosylate group, ester group with a nitrogen, oxygen, or carbon atom.

15

142. The method of synthesis of the compound of claim 131 wherein attaching the protected aminophthalhydrazide through the molecular linker to one of the aryl groups of diarylketene to form the precursor aminophthalimide-linked diarylketene is by a reaction of a nucleophilic group of the linker or aryl group of diarylketene with a leaving group of the linker or aryl group of diarylketene.

20

143. The method of synthesis of the compound of claim 142 wherein a linker is attached to the aryl group of diarylketene by reaction to form a bond between at least one of a nitrogen, oxygen, or carbon atom of the linker and at least one of a nitrogen, oxygen, or carbon atom of group of the protected aminophthalhydrazide by an addition or a substitution reaction of a leaving group.

25

144. The method of synthesis of the compound of claim 143 wherein a linker is attached to the aryl group of diarylketene by a substitution reaction of at least one of a halogen, tosylate group, ester group with a nitrogen, oxygen, or carbon atom.

30

145. The method of synthesis of the compound of claim 131 wherein the precursor aminophthalimide-linked diarylketene is further reacted by condensation with an aryl alkene aldehyde in a nonaqueous solvent, containing an acid catalyst to form B linked to the A precursor.

35

146. The method of synthesis of the compound of claim 145 wherein the precursor aminophthalimide-linked diarylketene is an aminophthalimide-substituted 1,1-diarylethene, the aryl alkene aldehyde is a p-aminophenyl alkene aldehyde such as p-(dimethylamino)-


cinnamaldehyde,

the nonaqueous solvent is acetic anhydride,

the acid catalyst is at least one of perchloric acid and tetrafluoroboric acid, and

the B linked to the A precursor comprises a aminophthalimide-substituted
5 multiarylpolymethine dye.

147. The method of synthesis of the compound of claim 145 wherein the precursor aminophthalimide-linked diarylketene comprises at least one of the formula



3a: R = N(CH₃)₂


3b: R = H

3c: R = OCH₃

3d: R = O(CH₂)₃CH₃


3e: R = (CH₂)₃CH₃

10 the aryl alkene aldehyde has the formula



4-(Dimethylamino)cinnamaldehyde, and

the precursor of A-B comprises at least one of the formula



22a: R = N(CH₃)₂

22b: R = H

22c: R = OCH₃

22d: R = O(CH₂)₃CH₃

22e: R = (CH₂)₃CH₃

15

148. The method of synthesis of the compound of claim 145 wherein the phthalimide moiety of the A precursor is converted to the phthalhydrazide A functionality by treating with hydrazine, forming A-B.

149. The method of synthesis of the compound of claims 148 wherein the B functionality is protected by reacting with an anion such as hydroxide, methoxide and amine, the A-B precursor is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent in inert atmosphere and then treated with acid such as perchloric acid, tetrafluoroboric acid to form A-B.

150. The method of synthesis of the compound of claim 131 wherein the phthalimide moiety of the A precursor of the precursor aminophthalimide-linked diarylketene is converted to the phthalhydrazide A functionality by treating with hydrazine, forming A attached to a B precursor.

151. The method of synthesis of the compound of claim 150 wherein the A-linked diarylketene is further reacted by condensation with an aryl alkene aldehyde in a nonaqueous solvent, containing an acid catalyst to form A-B.

152. The method of synthesis of the compound of claim 151 wherein the A-linked diarylketene is an aminophthalhydrazide-substituted 1,1-diarylethene,


the aryl alkene aldehyde is a p-aminophenyl alkene aldehyde such as p-(dimethylamino)-cinnamaldehyde,

the nonaqueous solvent is acetic anhydride,

the acid catalyst is at least one of perchloric acid and tetrafluoroboric acid, and

A-B comprises a aminophthalhydrazide-substituted multiarylpolymethine dye.

153. The method of synthesis of the compound of claim 152 wherein the A-linked diarylketene comprises at least one of the formula



3a: R = N(CH₃)₂


3b: R = H

3c: R = OCH₃

3d: R = O(CH₂)₃CH₃


3e: R = (CH₂)₃CH₃

the aryl alkene aldehyde has the formula



4-(Dimethylamino)cinnamaldehyde, and


A-B comprises at least one of the formula




5

154. The method of synthesis of the compound of claims 149 and 151 further comprising the step of reacting the B functionality with one nucleophilic species of a C functionality such as Foscarnet to form A-B-C.


10 155. The method of synthesis of the compound of claim 131 comprising the general steps given by following representative formula



- 3a:** R = N(CH₃)₂
3b: R = H
3c: R = OCH₃
3d: R = O(CH₂)₃CH₃
3e: R = (CH₂)₃CH₃






- 22a:** R = N(CH₃)₂
22b: R = H
22c: R = OCH₃
22d: R = O(CH₂)₃CH₃
22e: R = (CH₂)₃CH₃




- 23a:** R = N(CH₃)₂
23b: R = H
23c: R = OCH₃
23d: R = O(CH₂)₃CH₃
23e: R = (CH₂)₃CH₃



- 24a:** R = N(CH₃)₂
24b: R = H
24c: R = OCH₃
24d: R = O(CH₂)₃CH₃
24e: R = (CH₂)₃CH₃



156. The method of synthesis of the compound of claim 131 comprising the general steps given by following representative formula



- 3a:** $\text{R} = \text{N}(\text{CH}_3)_2$
3b: $\text{R} = \text{H}$
3c: $\text{R} = \text{OCH}_3$
3d: $\text{R} = \text{O}(\text{CH}_2)_3\text{CH}_3$
3e: $\text{R} = (\text{CH}_2)_3\text{CH}_3$




- 23a:** $\text{R} = \text{N}(\text{CH}_3)_2$
23b: $\text{R} = \text{H}$
23c: $\text{R} = \text{OCH}_3$
23d: $\text{R} = \text{O}(\text{CH}_2)_3\text{CH}_3$
23e: $\text{R} = (\text{CH}_2)_3\text{CH}_3$

- 24a:** $\text{R} = \text{N}(\text{CH}_3)_2$
24b: $\text{R} = \text{H}$
24c: $\text{R} = \text{OCH}_3$
24d: $\text{R} = \text{O}(\text{CH}_2)_3\text{CH}_3$
24e: $\text{R} = (\text{CH}_2)_3\text{CH}_3$

condensing the diarylketene with an aryl alkene aldehyde to form B
forming a protected aminophthalhydrazide such as aminophthalimide or aminophthalic acid diester,
adding a hydrocarbon linker to the protected aminophthalhydrazide, and
5 attaching the protected aminophthalhydrazide through the molecular linker to the aryl groups of B to form the precursor aminophthalimide-linked B, and
forming the A functionality from the precursor to form A-B.

158. The method of synthesis of the compound of claim 157 wherein at least one of the diaryl ketone and diarylketene is halo-substituted and the protected aminophthalhydrazide is attached through the linker by an amination reaction.

159. The method of synthesis of the compound of claim 158 wherein the halo-substituted diarylketene precursor compounds comprises the formula of at least one of



2a: R = N(CH₃)₂

2b: R = H

2c: R = OCH₃


2d: R = O(CH₂)₃CH₃

15 2e: R = (CH₂)₃CH₃ , and

the halo-substituted multiarylpolymethine dyes, such as 1-(p-bromophenyl)-1,5-bis(p-dimethylaminophenyl)-pentadienium perchlorate, are prepared by condensation with a p-aminophenyl alkene aldehyde such as p-(dimethylamino)cinnamaldehyde.

20 160. The method of synthesis of the compound of claim 158 wherein B is protected by reacting with an anion such as alkoxide and then coupled with A by amination of aryl halide such as the palladium-catalyzed amination of aryl halide to obtain the alkoxide-protected aminophthalimide-substituted multiarylpolymethine dye.


25 161. The method of synthesis of the compound of claim 160 wherein the protected aminophthalhydrazide-linked to B from the alkoxide-protected aminophthalimide-substituted multiarylpolymethine dye comprises at least one of the formula



- 22a:** R = N(CH₃)₂
22b: R = H
22c: R = OCH₃
22d: R = O(CH₂)₃CH₃
22e: R = (CH₂)₃CH₃

162. The method of synthesis of the compound of claim 160 wherein the alkoxide-protected aminophthalimide-substituted multiarylpolymethine dye is refluxed with hydrazine in a suitable solvent such as an alcoholic solvent to convert the amino-phthalimide moiety to the aminophthalhydrazide moiety and then treated with acid to generate A-B.

163. The method of synthesis of the compound of claim 162 wherein A-B comprises at least one of the formula



- 23a:** R = N(CH₃)₂
23b: R = H
23c: R = OCH₃
23d: R = O(CH₂)₃CH₃
23e: R = (CH₂)₃CH₃

164. The method of synthesis of the compound of claim 162 further comprising the step of reacting the B functionality with one nucleophilic species of a C functionality such as Foscamet to form A-B-C.

165. The method of synthesis of the compound of claim 157 wherein at least one of the diaryl ketone and diarylketene is halo-substituted and an aminophthalhydrazide is attached through the linker by an amination reaction.

166. The method of synthesis of the compound of claim 1 wherein the A functionality comprises an active oxalate and the B functionality comprises a multiarylpolymethine

photochromic dye wherein A is attached to aryl groups of B comprising the steps of
forming a halo-substituted diaryl ketone,

forming a halo-substituted diaryl ketene from the diaryl ketone,

amination of the halo-substituted diaryl ketene to give amino diarylketene,

5 substitution at the amino group of the ketene to forming the corresponding sulfonamide,
condensing the sulfonamide with a catalyst, and

react with oxalyl halide to form A-B.

10 167. The method of synthesis of the compound of claim 1 wherein the A functionality
comprises an cyclized active oxalate and the B functionality comprises a multiarylpolymethine
photochromic dye wherein A is attached to aryl groups of B comprising the steps of

forming a halo-substituted diaryl ketone,

forming a halo-substituted diaryl ketene from the diaryl ketone,

amination of the halo-substituted diaryl ketene to give amino diarylketene,

15 substitution at the amino group of the ketene to forming the corresponding sulfonamide,
reacting 2 molar proportions of a *N*-substituted aminodiarylketene with 1 molar oxalyl

halide to yield the *N,N'*-bisaryl oxamide,

condensing the oxamide with a catalyst to form A-B.




20 168. The method of synthesis of the compound of claim 167 wherein the halo-substituted
diaryl ketene is aminated using methods such as the palladium-catalyzed amination of aryl halide
with benzophenoneimine to give the amino diarylketene.

25 169. The method of synthesis of the compound of claim 167 wherein the amino groups of the
ketene are substituted forming the corresponding sulfonamide by reacting with sulfonyl
anhydride.

30 170. The method of synthesis of the compound of claim 167 wherein the oxamide is
condensed with an orthoester such as triethylorthoformate in a nonaqueous solvent such as acetic
anhydride containing acid catalyst such as tetrafluoroboric acid, to form the cyclized oxamido-
tetraarylpolymethine dye comprising A-B.

35 171. The method of synthesis of the compound of claim 166 further comprising the step of
reacting the B functionality with one nucleophilic species of a C functionality such as Foscarnet
to form A-B-C.

172. The method of synthesis of the compound of claim 167 wherein the general steps are
given by following representative formula



173. The method of synthesis of the compound of claim 1 wherein the A functionality comprises an active oxalate and the B functionality comprises a multiarylpolymethine photochromic dye wherein A is attached to aryl groups of B through a molecular linker

5 comprising the steps of

forming B comprising a functionalized tetraarylpolymethine dye,

reacting a substituted amine with a sulfonyl anhydride to form a substituted alkyl

sulfonamide,

reacting the substituted alkyl sulfonamide with an oxalyl derivative to form a substituted oxamide,

reacting the substituted oxamide with the functionalized tetraarylpolymethine dye to form

5 A-B comprising a cyclized oxamido-tetraarylpolymethine.

174. The method of synthesis of the compound of claim 173 wherein the substituted amine is N-2-bromoethylsulfamide.

10 175. The method of synthesis of the compound of claim 173 wherein the oxalyl derivative is oxalyl chloride.

176. The method of synthesis of the compound of claim 173 wherein the oxamide is a N-2-bromoethyl-N-sulfonyloxamide derivative.

15

177. The method of synthesis of the compound of claim 173 wherein the oxalyl derivative is oxalyl chloride.






178. The method of synthesis of the compound of claim 173 wherein the functionalized
20 tetraarylpolymethine derivative is a salt of a 1,5-bis(4-hydroxyphenyl)-1,5-diarylpentadiene derivative.





179. The method of synthesis of the compound of claim 173 wherein the cyclized oxamido-
tetraarylpolymethine A-B compound is a 1,5-(4,4'-(2,2'-N,N'-
25 disulfonyloxamidodiethoxy)phenyl)-1,5-diarylpentadiene cation derivative.

180. The method of synthesis of the compound of claim 173 further comprising the step of
reacting the B functionality with one nucleophilic species of a C functionality such as Foscarnet
to form A-B-C.

30

181. The method of synthesis of the compound of claim 173 comprising the general steps
given by following representative formula





182. The method of synthesis of the compound of claims 1, 70, 73, 76, 105, 131, 157, 166, 167, 173 wherein one or more of the moieties are modified to further candidate components by addition of functional groups.

5

183. The method of synthesis of the compound of claim 182 wherein each of A, B, and C are

modified to further candidate components by addition of functional groups one the group comprising alkyl, cycloalkyl, alkoxycarbonyl, cyano, carbamoyl, heterocyclic rings containing C, O, N, S, sulfo, sulfamoyl, alkoxysulfonyl, phosphono, hydroxyl, halogen, alkoxy, alkylthiol, acyloxy, aryl, alkenyl, aliphatic, acyl, carboxyl, amino, cyanoalkoxy, diazonium, carboxyalkylcarboxamido, alkenylthio, cyanoalkoxycarbonyl, carbamoylalkoxycarbonyl, alkoxy carbonylamino, cyanoalkylamino, alkoxycarbonylalkylamino, sulfoalkylamino, alkylsulfamoylalkylamino, oxido, hydroxy alkyl, carboxy alkylcarbonyloxy, cyanoalkyl, carboxyalkylthio, arylamino, heteroarylamino, alkoxycarbonyl, alkylcarbonyloxy, cyanoalkoxy, alkoxycarbonylalkoxy, carbamoylalkoxy, carbamoylalkyl carbonyloxy, sulfoalkoxy, nitro, alkoxyaryl, halogenaryl, amino aryl, alkylaminoaryl, tolyl, alkenylaryl, allylaryl, alkenyloxyaryl, allyloxyaryl, cyanoaryl, carbamoylaryl, carboxyaryl, alkoxycarbonylaryl, alkylcarbonyoxyaryl, sulfoaryl, alkoxysulfoaryl, sulfamoylaryl, and nitroaryl.

184. The compound comprising the formula A'-B-C
where the A' is a chemiluminescent moiety precursor,
B is an energy acceptor moiety, and
C is a biologically active moiety.

185. The compound comprising the formula A'-B
where the A' is a chemiluminescent moiety precursor, and
B is an energy acceptor moiety.


186. The compound comprising the formula A'-B'
where the A' is a chemiluminescent moiety precursor, and
B' is an energy acceptor moiety precursor.

187. The compound of claims 184, 185, and 186 wherein A' is a precursor to generate a phthalhydrazide.


188. The compound of claim 187 wherein A' is at least one of phthalimide, aminophthalic acid diester, aminophthalic acid dihydrazide, and aminophthalic anhydride.

189. The compound of claim 187 wherein A' is a phthalhydrazide protected by a hydrolyzable group.

190. The compound of claims 184 and 185 wherein the structure is given by at least one of the general formula



and



wherein the functionality **A'** is at least one of precursor of an aminophthalhydrazide derivatives, sulfonyloxamides and active oxalates,

the functionality **B** is at least one of 1,1,5,5-tetrakisarylpentadiene derivative,


the functionality **C** is a drug molecule such as Foscarnate, or ddc, and

R is a functional group, and


L is a linker such as an aliphatic chain between **A'** and **B**.

191. The compound of claim 190 wherein the **L** functionality is between one 20 carbon atoms.

192. The compound of claims 184 and 185 wherein the structure is given by at least one of the general formula



and



5

wherein the functionality **A'** is at least one of precursor of an aminophthalhydrazide derivatives, sulfonyloxamides and active oxalates,

the functionality **B** is a 1,1,5-trisubstituted pentadiene derivatives,

the functionality **C** is a drug molecule such as Fosfarnate, or ddc, and


10





R is a functional group, and

L is a linker such as an aliphatic chain between **A'** and **B**.


193. The compound of claim 192 wherein the **L** functionality is between one 20 carbon atoms.

15 194. The compound of claims 184, 185, 186, 190, and 192 wherein the structure of **A'** is given by at least one of the general formula





5 , and




wherein R^1 , R^2 , R^3 , R^4 are one of the following groups: alkyl, alkoxy, alkylamino or hydrogen, and R, R', and R'' is at least one of alkyl, cycloalkyl, alkoxy, carbonyl, cyano, carbamoyl, heterocyclic rings containing C, O, N, S, sulfo, sulfamoyl, alkoxy, sulfonyl, phosphono, hydroxyl, halogen, alkoxy, alkylthiol, acyloxy, aryl, alkenyl, aliphatic, acyl, carboxyl, amino, cyanoalkoxy, diazonium, carboxyalkylcarboxamido, alkenylthio, cyanoalkoxy, carbamoylalkoxy, alkoxy carbonyl, cyanoalkylamino, alkoxy carbonylalkylamino, sulfoalkylamino, alkylsulfamoylalkylamino, oxido, hydroxy alkyl, carboxy alkylcarbonyloxy, cyanoalkyl, carboxyalkylthio, arylamino, heteroaryl, alkoxy carbonyl, alkylcarbonyloxy, cyanoalkoxy, alkoxy carbonylalkoxy, carbamoylalkoxy, carbamoylalkyl carbonyloxy, sulfoalkoxy, nitro, alkoxyaryl, halogenaryl, amino aryl, alkylaminoaryl, tolyl, alkenylaryl, allylaryl, alkenyloxyaryl, allyloxyaryl, cyanoaryl, carbamoylaryl, carboxyaryl, alkoxy carbonylaryl, alkylcarbonyloxyaryl, sulfoaryl, alkoxy sulfonyl, sulfamoylaryl, and nitroaryl.

195. The compound of claims 184, 185, 186, 190, and 192 wherein A' is a precursor to generate an active oxalate.


196. The compound of claim 195 wherein A' is a sulfonamide.

197. The compound of claim 196 wherein A' is a precursor to generate a cyclized active oxalate.
- 5 198. The compound of claim 196 wherein A' is a sulfonamide bound to at least one of a diaryl ketone, halo-substituted diaryl ketone, amino-substituted diaryl ketone, diaryl ketene, halo-substituted diaryl ketene, and amino-substituted diaryl ketene.
- 10 199. The compound of claims 185 and 186 wherein A'-B or A'-B' is a sulfonamide precursor for a cyclized sulfonyloxamide derivative bound to a photochromic dye comprising A-B.
200. The compound of claim 197 wherein A' is attached to aryl groups of B through a molecular linker.
- 15 201. The compound of claim 196 wherein A' is at least one of substituted alkyl sulfonamide and a substituted oxamide.
202. The compound of claim 201 wherein the substituted amide is a N-2-bromoethylsulfamide derivative.
- 20 203. The compound of claim 201 wherein the oxamide is a N-2-bromoethyl-N-sulfonyloxamide derivative.
- 25 204. The compound of claims 184, 190, and 192 wherein B comprises a functionalized tetraarylpolymethine derivative such as a salt of a 1,5-bis(4-hydroxyphenyl)-1,5-diarylpentadiene derivative.
- 30 205. The compound of claim 199 wherein the cyclized oxamido-tetraarylpolymethine A-B compound is a 1,5-(4,4'-(2,2'-N,N'-disulfonyloxamidodiethoxy)phenyl)-1,5-diarylpentadiene cation derivative.
206. The compound of claim 205 given by the general formula




207. The compound of claim 184 comprising A-B-C wherein the cyclized oxamido-tetraarylpolymethine A-B compound is a 1,5-(N,N'-disulfonyloxamido)phenyl-1,5-diarylpentadiene cation; or a 1,5-(4,4'-(2,2'-N,N'-disulfonyloxamidodiethoxy)phenyl-1,5-diarylpentadiene cation derivative and C is abiological active moiety such as the drug Foscarnet.

208. The compound of claim 207 given by the general formula



10 wherein L is a linker with 0 to 20 chain atoms.


209. The compound of claims 184, 185, 186, 190, and 192 wherein the structure of A' is a oxalylate given by the general formula



15
20 wherein X is O or N and y is O or N; R's are one the following groups: hydrogen, alkyl, cycloalkyl, alkoxy carbonyl, cyano, carbamoyl, heterocyclic rings containing C, O, N, S, sulfo, sulfamoyl, alkoxy sulfonyl, phosphono, hydroxyl, halogen, alkoxy, alkylthiol, acyloxy, aryl, alkenyl, aliphatic, acyl, carboxyl, amino, cyanoalkoxy, diazonium, carboxyalkylcarboxamido, alkenylthio, cyanoalkoxy carbonyl, carbamoylalkoxy carbonyl, alkoxy carbonylamino,


cyanoalkylamino, alkoxycarbonylalkylamino, sulfoalkylamino, alkylsulfamoylalkylamino, oxido, hydroxy alkyl, carboxy alkylcarbonyloxy, cyanoalkyl, carboxyalkylthio, arylamino, heteroaryl, alkoxycarbonyl, alkylcarbonyloxy, cyanoalkoxy, alkoxycarbonylalkoxy, carbamoylalkoxy, carbamoylalkyl carbonyloxy, sulfoalkoxy, nitro, alkoxyaryl, halogenaryl, amino aryl, alkylaminoaryl, tolyl, alkenylaryl, allylaryl, alkenyloxyaryl, allyloxyaryl, cyanoaryl, carbamoylaryl, carboxyaryl, alkoxycarbonylaryl, alkylcarbonyloxyaryl, sulfoaryl, alkoxysulfoaryl, sulfamoylaryl, and nitroaryl.

210. The compound of claim 209 wherein A' is a sulfonyl oxamide given by the general formula




wherein L is aliphatic chain linker with 0 to 20 chain atoms.

211. The compound of claims 184, 185, 186, 190, and 192 wherein the structure of A' is given by the general formula




where X is a leaving group such as halide.


212. The compound of claim 184 and 190 wherein the structure of A'-B is given by the general formula




213. The compound of claim 186 wherein the structure of A'-B' is given by the general formula



A-B' is given by the general formula



- 5 214. The compound comprising the structure of A-B-C given by the general formula



- 10 215. The compound of claim 186 wherein the B' moiety is a precursor photochromic compound.

216. The compound of claims 184, 185, 190, 192, and 215 wherein the photochromic compound comprises one which demonstrate photochromic behavior with electromagnetic radiation and bleaching agents.

15

217. The compound of claim 216 wherein the photochromic compound comprises a cationic dye.

218. The compound of claim 217 wherein the cationic dye comprises at least one of a di and

triarylmethane dyes, triarylmethane lactones and cyclic ether dyes, cationic indoles, pyronines, phthaleins, oxazines, thiazines, acridines, phenazines, and anthocyanidins, and cationic polymethine dyes and azo and diazopolymethines, styryls, cyanines, hemicyanines, dialkylaminopolyenes, and other related dyes.

5


219. The compound of claim 218 wherein the cationic dye comprises at least one of

	Malachite Green	42000
	Helvetia Green	42020
10	Basic Blue 1	42025
	Brilliant Blue	
	Setoglaurine	
	Basic Green 1	42040
	Brilliant Green	
15	Acid Blue 1	42045
	Xylene Blue VS	
	Patent Blue V	
	Alphazurine 2G	
	Acid Blue 3	42051
20	Brilliant Blue V	
	Patent Blue V	
	Food Green 3	42053
	FDC Green 3	
	Acid Green 6	42075
25	Light Green SF Bluish	
	Acid Blue 7	42080
	Xylene Blue AS	
	Patent Blue A	
	Acid Green 3	42085
30	Acid Blue 9	42090
	Erioglaurine	
	Acid Green 5	42095
	Light Green SF Yellowish	
	Acid Green 9	42100
35	Erioviridene B	
	Acid Blue 147	42135
	Xylene Cyanol FF	
	Basic Red 9	42500
	Pararosaniline	


	Basic Violet 14	42510
	Fuchsin	
	Magenta	
	Basic Fuchsin	42510B
5	Basic Violet 2	42520
	New Magenta	
	Hoffman Violet	42530
	Iodine Violet	
	Basic Violet 1	42535
10	Methyl Violet	
	Basic Violet 13	42536
	Methyl Violet 6B	
	Basic Violet 3	42555
	Crystal Violet	
15	Gentian Violet	
	Iodine Green	42556
	Basic Blue 8	42563
	Victoria Blue 4R	
	Acid Blue 13	42571
20	Fast Acid Violet 10B	
	Acid Blue 75	42576
	Eriocyanine A	
	Methyl Green	42585
	Ethyl Green	42590
25	Basic Violet 4	42600
	Ethyl Violet	
	Acid Violet 49	42640
	Wool Violet 5BN	
	Acid Blue 15	42645
30	Brilliant Milling Blue B	
	Acid Violet 17	42650
	Acid Violet 6B	
	Wood Violet 4BN	
	Formyl Violet	
35	Acid Violet 5BS Conc.	
	Acid Violet 19	42685
	Acid Fuchsin	
	Red Violet 5R	42690
	Acid Blue 22	42755

	Aniline Blue	
	Soluble Blue	
	Solvent Blue 3	42775
	Solvent Blue 3	42780
5	Methyl Blue	
	Aurin	43800
	Mordant Blue 3	43820
	Eriochrome Cyanine R	
	Acid Green 16	44025
10	Naphthalene Green V	
	Pontacyl Green NV Extra	
	Basic Blue 11	44040
	Victoria Blue R	
	Basic Blue 15	44085
15	Night Blue	
	Acid Green 50	44090
	Wool Green S	
	Kiton Green S. Conc.	
	Basic Green 3	
20	Sevron Green B	
	Brilliant Blue F & R Extra	
	Brilliant Green Sulfonate	
	Hexakis (hydroxyethyl)	


Pararosanine




New Green




Phenolphthalein




Malachite Green Ethiodide




Hydroxyalkylated Pararosanines




Hydroxyalkylated New Fuchsin




New Yellow




Doebner's Violet




New Red




Bis(hydroxyethyl) Doebner's Violet




"New Magenta"










Tetrakis(hydroxyethyl) Doebner's Violet




Trichloro Crystal Violet



Slow Red





5

^a Only the cyanide, bisulfite, and hydroxide ions are considered, regardless of the other anions present in the solution.

10 ^b More detailed descriptions of the compositions of photochromic materials tested are given in Macnair's review [255; tables 1A-4].

^c Ethanol.

^d Diethyl ether.

^e 1,2-Dichloroethane.

15 ^f 1,1-Dichloroethane, cyclohexane-1,1-dichloroethane, or cyclohexane-1,2-dichloroethane mixtures.

^g Benzene.

^h Dimethylsulfoxide, neat and aqueous.

- ⁱ Acetone.
- ^j **ACETIC ACID.**
- ^k Ethyl acetate.
- ^l Ethyl bromide.
- 5 ^m 2-Methoxyethanol.
- ⁿ Chloroform.
- ^o Ethanol with KCN.
- ^p Ethanol with KOH.
- ^q Carboxylic acids-acetic to stearic; hydrocinnamic acid; ethyl and butyl acid phthalates.
- 10 ^r Octadecylnitrile, tributyl phosphate, aniline, 2-(p-tert-butylphenoxy)ethanol, tetraethyleneglycol dimethyl ether, or poly(ethylene glycols).
- ^s Amides-formamide to stearamide; methylformamide or methylacetamide; dimethyl- or diethyl-formamide or acetamide.
- ^t Three-to-one solutions of cellulose acetate with any of the following five-to-one plasticizer mixtures:
- 15 butyl stearate, Polyethylene Glycol 600-butyl acetoxystearate, butyl stearate, or Dowanol EP-butyl acetoxystearate.
- ^u Water containing SO₂.
- ^v Water containing bisulfite and papain.
- ^w Poly(vinyl alcohol) with dimethylsulfoxide (5:1).
- 20 ^x Films, containing residual solvent, cast from the following solutions: ethanol-acetone solutions of vinyl acetate-vinyl alcohol copolymer; aqueous poly(vinyl alcohol); aqueous poly(vinyl pyrrolidone); or aqueous methyl vinyl ether-maleic acid copolymer.
- ^y Methanol-dioxane with aqueous NH₄HSO₃.
- ^z Paper impregnated with a toluene solution of poly(methyl methacrylate), stearic acid, and 2-(p-tert-butylphenoxy)ethanol, then dried.
- 25 ^{aa} Intramicellar impregnation of cellulose with the following swelling agents: n-propylamine, n-butylamine, n-hexylamine, 2-aminoethanol, dimethylformamide, acetic acid, dimethylsulfoxide, methylacetamide, dimethylacetamide, or formamide.
- ^{bb} Films cast from an approximately 4:3 mixture of a 20% solution and cellulose acetate butyrate in toluene-ethyl acetate (1:1) and triallylcyanurate in dioxane.
- 30 ^{cc} **FILMS CAST FROM A 2:1 MIXTURE OF A 25% SOLUTION OF CELLULOSE ACETATE BUTYRATE IN TOLUENE-ETHYL ACETATE (1:1) AND THE TITANIUM ESTERS OF N,N,N', N'-TETRAKIS(2-HYDROXYPROPYL) ETHYLENEDIAMINE.**
- ^{dd} Pure water.
- 35 ^{ee} Films cast from aqueous gelatin or other hydrocolloids.
- ^{ff} Dimethylsulfoxide with methanolic KCN.
- ^{gg} 2-Methoxyethanol with methanolic KCN.
- ^{hh} Water or aqueous methanol containing bisulfite.
- ⁱⁱ Paper impregnated with m-dimethoxybenzene, acetonitrile, acetic acid, or phenyl methyl carbinol.

ij Ethanol-benzene.

kk Aqueous ethanol, methanol, aqueous methanol, aqueous acetone, benzene-methanol, carbon tetrachloride-methanol, cyclohexane-methanol, or chloroform-methanol.

ll Films cast from 3:1 solutions of cellulose acetate and either Polyethylene Glycol 600 .RTM. or ethylene glycol phenyl ether as plasticizer.

mm Films, containing residual solvent, cast from solutions of either cellulose acetate in 2-methoxyethanol or poly(vinyl alcohol) in aqueous ethanol.

nn Films, containing residual solvent, cast from solutions of either cellulose acetate butyrate in 2-methoxyethanol or poly(vinyl acetate) in methanol.

oo Ethanol containing ammonia.

pp Aqueous methanol containing $\text{NH}_4 \text{HSO}_3$ and urease.

qq Aqueous methanol containing $\text{NH}_4 \text{HSO}_3$, with or without sodium dithionite.

rr Aqueous acid at pH 1.

ss Aqueous ammonia containing KCN.


tt Paper impregnated with aqueous solutions with or without hydrocolloids.





uu 2-Methoxyethanol containing HCl.


vv Aqueous methanol containing $\text{NH}_4 \text{HSO}_3$, and glucose oxidase.

ww 9:1 Methanol-water.


xx Aqueous NaOH.















	Ar	n
5	C ₆ H ₅	0, 1, 2
	4-(CH ₃) ₂ NC ₆ H ₄	0, 1, 2
	4-(CH ₃) ₂ CHC ₆ H ₄	0, 1, 2, 3, 4
	4-CH ₃ OC ₆ H ₄	0, 1, 2
	4-C ₄ H ₉ OC ₆ H ₄	0, 1, 2
10	3-CH ₃ C ₆ H ₄	1, 2
	4-t-C ₄ H ₉ C ₆ H ₄	1, 2
	4-C ₂ H ₅ OC ₆ H ₄	1, 2
	4-C ₅ H ₁₁ C ₆ H ₄	1, 2
	4-FC ₆ H ₄	1
15	4-Fsub ₃ CC ₆ H ₄	1
	2-(C ₆ H ₅) ₂ NC ₆ H ₄	1
	3,4-H ₂ N(OCH ₃)C ₆ H ₃	1
	2-Naphthyl	1, 2
	4-ClC ₆ H ₄	2
20	2,4-Cl ₂ C ₆ H ₃	2
	1-Naphthyl	2










25	R	R
----	---	---













5



Miscellaneous polyenes








Basic Red 13












Basic Violet 7





Basic Red 14
Basic Red 15
Basic Violet 15






























Salt-isomerism type phototropic dyes


Night Blue




Victoria Blue R




Brilliant Milling Blue B
Brilliant Blue F & R Ex.
Eriocyanine A




Methyl Blue




Aniline Blue




Eriochrome Cyanine R




Methyl Violet 6B




Iodine Green




Aniline Blue




Wool Violet 5 BN




Wool Violet 4 EM




Light Green SF
Yellowish




Iodine Violet




Methyl Violet




Crystal Violet




Red-Violet 5R




Brilliant Green "B"




Di-[4(N,N-diethylamino)phenyl]-[4(N,N-diethylamino-2-methyl) phenyl] methyl carbonium




Tri-[4(N,N-dipropylamino)phenyl] methyl carbonium




Di-[4(N,N-diethylamino)phenyl]-[4(ethylamino)-phenyl] methyl carbonium




Di-[4(N,N-diethylamino)phenyl]-[4(N,N-diethylamino)naphthyl] methyl carbonium



Di-[4(N,N-dimethylamino)phenyl]-[4(hydroxy)phenyl]
methyl carbonium



Tri-[4(N-propylamino)phenyl] methyl carbonium




Hectolene Blue DS-1398


Hectolene Blue DS-1823

Sevron Brilliant Red 4G

Di-[4(N,N-dimethylamino)phenyl]-[4(hydroxy)phenyl]
methyl carbonium



Tri-[4(N-propylamino)phenyl] methyl carbonium



Hectolene Blue DS-1398

Hectolene Blue DS-1823

Sevron Brilliant Red 4G


Genacryl Red 6B

Genacryl Pink G


Sevron Brilliant - Red B

Sevron Brilliant - Red 3B


1,5-bis-[4(N,N-dimethylamino)phenyl]-1,5-bis-
(phenyl)divinyl carbonium trifluoroacetate




1,1,3,3-tetrakis[4(N,N-dimethylamino)phenyl]
vinyl carbonium perchlorate




1,5-bis-[4(N,N-dimethylamino)phenyl]-1,5-bis-
(phenyl) divinyl carbonium *p*-toluenesulfonate




1,7-bis[4(N,N-dimethylamino)phenyl]-1,7-bis-
(2,4-dichlorophenyl) trivinyl carbonium perchlorate




Di-[4(N,N-dimethylamino)phenyl vinyl]-[2,4-di-
phenyl-6-methane thiopyran] methyl carbonium perchlorate




1,7-bis-[4(N,N-dimethylamino)phenyl]-1,7-bis-
(4-chlorophenyl) trivinyl carbonium trifluoroacetate




1,1,3-tris-[4-(N,N-dimethylamino)phenyl] divinyl
carbonium perchlorate




1,1,7,7-tetrakis-[4-(N,N-dimethylamino)phenyl]
trivinyl carbonium perchlorate




1,3-bis-[4-(N,N-dimethylamino)phenyl]-1,3-bis-
(phenyl) vinyl carbonium perchlorate




1,1,5,5-tetrakis-[4-(N,N-dimethylamino)phenyl]
divinyl carbonium perchlorate




1,5-bis-[4-(N,N-dimethylamino)phenyl]-1,5-bis-
(phenyl) divinyl carbonium perchlorate




1,7-bis-[4-(N,N-dimethylamino)phenyl]-1,7-bis-(phenyl) trivinyl carbonium trifluoroacetate




1(1,3,3-trimethyl indoline)-2-[4-(N,N-dimethylamino)phenyl] ethylene carbonium perchlorate




1(1,3,3-trimethyl indoline)-4-[4-(N,N-dimethylamino)phenyl] butylene carbonium perchlorate




1,1,3,3-tetrakis-[4-(N,N-diethylamino)phenyl] vinyl carbonium perchlorate




1,1-bis-[4-(N,N-diethylamino)phenyl]-3,3-bis-[4-(N,N-dimethylamino)phenyl] vinyl carbonium perchlorate




1,1,5,5-tetrakis-[4-(N,N-diethylamino)phenyl]
divinyl carbonium perchlorate




1,1-bis-[4-(N,N-dimethylamino)phenyl]-3-[4-(amino)
phenyl]-3-methylvinyl carbonium perchlorate




Tris-[1,1-bis-[4-(N,N-dimethylamino)phenyl]
ethylene] methyl carbonium perchlorate




Tris-[1,1-bis-[4-(N,N-diethylamino)phenyl]
ethylene] methyl carbonium perchlorate




1,1,5-tris-[4-(N,N-dimethylamino)phenyl] divinyl
carbonium perchlorate




N[4-(N,N-dimethylamino) cinnamylidene] auramine




1,1-bis-[4-(N,N-dimethylamino)phenyl]-3,4-bis-
(phenyl))-3,4-diazo butene carbonium



1,1,5,5-tetrakis-[4-(N,N-dimethylamino)phenyl]-
2,3-diazo pentene carbonium





N-(N',N'-dimethylamino cinnamylidene)-N,N-diphenyl
ammonium



Azo Polymethines

Dyes of the general structural type






5 220. The compound of claim 186 wherein B' comprises a diaryl ketone.

221. The compound of claim 186 wherein B' comprises a diaryl ketene.

222. A compound of claim 220 wherein the diaryl ketone comprises derivatives of the
10 following representative formula




3a: R = N(CH₃)₂

3b: R = H


3c: R = OCH₃

3d: R = O(CH₂)₃CH₃

3e: R = (CH₂)₃CH₃



3f




12a: R=OCH₃


12b: R=O(CH₂)₃CH₃

12c: R=(CH₂)₃CH₃

5



12d




12a: R=OCH₃

12b: R=O(CH₂)₃CH₃

12c: R=(CH₂)₃CH₃

12d: R=N(CH₃)₂

- 10 223. A compound of claim 221 wherein the diaryl ketene comprises derivatives of the following representative formula




3a: $R = N(CH_3)_2$

3b: $R = H$


3c: $R = OCH_3$

3d: $R = O(CH_2)_3CH_3$

3e: $R = (CH_2)_3CH_3$



3f




12a: $R = OCH_3$


12b: $R = O(CH_2)_3CH_3$

12c: $R = (CH_2)_3CH_3$

5



12d




12a: $R = OCH_3$

12b: $R = O(CH_2)_3CH_3$

12c: $R = (CH_2)_3CH_3$

12d: $R = N(CH_3)_2$

224. A compound of claims 184, 185, 190, 192, and 215 wherein A'-B comprises derivatives of the following representative formula



4a: $R = N(CH_3)_2$, $X = ClO_4^-$


4b: $R = H$, $X = BF_4^-$

4c: $R = OCH_3$, $X = BF_4^-$

4d: $R = O(CH_2)_3CH_3$, $X = BF_4^-$

4e: $R = (CH_2)_3CH_3$, $X = BF_4^-$

5




19a: $R = OCH_3$

19b: $R = O(CH_2)_3CH_3$

19c: $R = (CH_2)_3CH_3$

19d: $R = N(CH_3)_2$



22a: $R = N(CH_3)_2$


22b: $R = H$

22c: $R = OCH_3$

22d: $R = O(CH_2)_3CH_3$

22e: $R = (CH_2)_3CH_3$

10




22f


225. The compound of claims 184, 190, and 192 wherein the C moiety is at least one of the group of

5


Captopril




Prostaglandin E₂




2,3-dichloro- α -methylbenzylamine




3'-deoxy-S-adenosyl-L-homocysteine




Sinefungin




3,5-diiodo-4-hydroxybenzoic acid




6,6'-dithiobis (9-B-D-ribofuranosylpurine)




γ -aminobutyric acid




Gabaculine




N-(5'-phosphopyridoxy)-
4-aminobutyric acid




4-amino-hex-5-enoic acid




Baclofen




Adenosine




3-hydroxy-3-methyl-
glutarate




Campactin




But-3-ynoyl-CoA




Suramin




L-3-iodotyrosine




L-3-iodo-α-methyltyrosine




Disodium cromoglycate




Adenosine
3',5'-cyclic monophosphate




D,L-B-(5-hydroxy-3-indolyl)- α -hydrazino- α -propionic acid




D,L- α -hydrazino- α -methyl-dopa




α -methyl-dopa




5-(3,4-dihydroxycinnamoyl)salicylic acid




N-(phosphonacetyl)-L-aspartate




P-glycolohydroxamate




5-(p-sulfamylphenyl)-azosalicylic acid




Coformycin



Formycin B




Thioinosinate




Phosphonoformate




Phosphonoacetate




Ridavirin




Sotalol




Cimetidine




Fuscaric acid




2-mercaptoethylamine




Mimosine




U-7130




Iproniazid




Trans-4-aminoocrotonic acid




NSD 1055




Nicotinic acid




Kynurenic acid




Lentysine




Orotic acid




Polyoxin D



Cephalosporin



Penicillin



5

226. The compound of claims 184, 185, 186, 190, 192, 221, 213, 215, and 225 wherein one or more of the moieties or their precursors are modified to further candidate components by addition of functional groups.

- 10 227. The compound of claim 226 wherein each of at least A, B, C, A', B', and C' wherein C' is a modified biologically active compound or their precursors are modified to further candidate components by addition of functional groups one the group comprising alkyl, cycloalkyl, alkoxy carbonyl, cyano, carbamoyl, heterocyclic rings containing C, O, N, S, sulfo, sulfamoyl, alkoxy sulfonyl, phosphono, hydroxyl, halogen, alkoxy, alkylthiol, acyloxy, aryl, alkenyl, aliphatic, acyl, carboxyl, amino, cyanoalkoxy, diazonium, carboxyalkylcarboxamido, alkenylthio, 15 cyanoalkoxy carbonyl, carbamoylalkoxy carbonyl, alkoxy carbonylamino, cyanoalkylamino, alkoxy carbonylalkylamino, sulfoalkylamino, alkylsulfamoylalkylamino, oxido, hydroxy alkyl, carboxy alkylcarbonyloxy, cyanoalkyl, carboxyalkylthio, arylamino, heteroaryl amino, alkoxy carbonyl, alkylcarbonyloxy, cyanoalkoxy, alkoxy carbonylalkoxy, carbamoylalkoxy,

carbamoylethyl carbonyloxy, sulfoalkoxy, nitro, alkoxyaryl, halogenaryl, amino aryl, alkylaminoaryl, tolyl, alkenylaryl, allylaryl, alkenyloxyaryl, allyloxyaryl, cyanoaryl, carbamoylaryl, carboxyaryl, alkoxycarbonylaryl, alkylcarbonyoxyaryl, sulfoaryl, alkoxysulfoaryl, sulfamoylaryl, and nitroaryl.